

2022-2023

ERNSTSEED.COM

MEADVILLE, PENNSYLVANIA





EST. 1964

MEADVILLE,  
PENNSYLVANIA

IT IS THE MISSION OF ERNST CONSERVATION SEEDS TO MAKE AVAILABLE TO EASTERN NORTH AMERICA THE KEY NATIVE AND NATURALIZED SPECIES OF PLANTS FOR RESTORATION, RECLAMATION, CONSERVATION, WILDLIFE AND POLLINATOR HABITAT ENHANCEMENT, RENEWABLE BIOMASS ENERGY, AND BEAUTIFICATION OF OUR NATION.

*We will identify, collect, and propagate new species and ecotypes that meet our clients' needs, from eastern Canada to the southeastern United States. Our native seeds are produced from species considered to be the most significant foundation of an effective native restoration or reclamation project. To ensure that our customers receive a quality product, all our seeds are harvested, conditioned, and tested under the highest quality standards.*

*In addition to 10,000 acres of native seed production at Ernst, we also supply seed from some of North America's top producers and collectors.*

*Ernst Conservation Seeds and its suppliers collect, grow, and process all our products in an ecologically sustainable and renewable manner.*

## DIRECTIONS

### TO ERNST CONSERVATION SEEDS

(Note: Some internet and GPS directions are not accurate)

**Located in Northwest Pennsylvania, 80 miles north of Pittsburgh and 40 miles south of Lake Erie off I-79:**

Take I-79 to **Meadville Exit 147A** (US 322 East to Meadville).

Travel on **US 322 East** to 2nd traffic light.

Turn right onto **Mercer Pike** and travel approximately 1000' before turning left to continue on Mercer Pike (follow the sign to "Ernst Bike Trail").

Drive 2 miles on Mercer Pike.

After crossing the railroad tracks, turn left into the **Ernst lane**.



# CONTENTS

ORDERING INFORMATION	2	BIOMASS	60
LETTER FROM CALVIN ERNST	3	BIOENGINEERING MATERIALS	64
ERNST TEAM	4	GRASSES & GRASS-LIKE SPECIES	69
DEFINING NATIVE VS. NATURALIZED	6	HERBACEOUS FLOWERING SPECIES	85
UNDERSTANDING PURE LIVE SEED (PLS)	7	WOODY TREES, SHRUBS & VINES	109
EXPECTATIONS OF NATIVE SPECIES	8	FORAGE, COVER CROPS, WILDLIFE FOOD PLOTS & BIOFUELS	115
ESTABLISHMENT GUIDE	12	INDEX	122
OBSTACLES TO A SUCCESSFUL MEADOW ESTABLISHMENT	55		

## Planting Guides for Native & Naturalized Seed Projects

Upland & Meadow Sites	19
Wildlife Habitat & Food Plot Sites	23
Pollinator-Friendly Sites	26
Riparian Sites	29
Disturbed Sites & Steep Slopes	32
Utility Right-of-Way Sites	36
Wet Meadow & Wetland Sites	38
Partially Shaded Sites	41
Stormwater Management Sites	44
Southeastern U.S. Sites	48
Biomass Production Sites	52

*Aster pilosus* (Heath Aster)  
with a Monarch Butterfly.

## CONNECT:

  } @ErnstSeeds

  } Ernst Conservation Seeds

# ORDERING INFORMATION

## PLACING AN ORDER

Our hours are Monday through Friday, 8 a.m. to 5 p.m. ET



**TOLL FREE:**  
800-873-3321  
**PHONE:**  
814-336-2404



**E-MAIL:**  
sales@ernstseed.com



**FAX:**  
814-336-5191



**STREET/MAILING ADDRESS:**  
8884 Mercer Pike  
Meadville, PA 16335

## TERMS AND CONDITIONS

### ORDER MINIMUMS

- \$25.00 order minimum before freight and tax.
- Mixes must be ordered in whole pound increments.
- 5% custom mix charge.

**PAYMENT:** Visa, MasterCard, Discover, and American Express are accepted forms of payment. Checks and money orders are also acceptable with placement of order. Sorry, Ernst cannot process COD orders.

**FREIGHT:** Orders are shipped via UPS or commercial freight (as applicable) and require a street address. Freight is pre-paid and added to your invoice unless prior arrangements are made with an Ernst sales associate.

**OTHER FEES:** Other fees may include, but are not limited to, small order fees, pallet packaging, custom orders, and bioengineering surcharges. We may require payment in advance for certain custom or special orders.

**CANCELLATIONS & RETURNS:** We cannot guarantee that cancellation requests can be processed within 48 hours of

the shipping date. Cancellation requests are subject to Ernst approval.

All plants are guaranteed to be what is specified on the plant tag or shipping documents and healthy upon receipt. Since we cannot guarantee care of such items after they leave our dock, all live plant material claims must be made within 48 hours of receipt.

All orders are subject to acts of natural destruction crop failures; liability is limited to refund or replacement.

Returns must be approved by Ernst and are subject to a restocking fee. Individual items and Ernst stock mixes are subject to a 10% restocking fee and must be returned within 30 days in the original packaging. Custom orders cannot be returned once the individual items are mixed. Bioengineering order cancellations are subject to our approval and assessed a 25% cancellation fee. Bioengineering orders are non-returnable.

**Prices are subject to change without notice.  
Please call for pricing and availability.**



Ernst Seeds Headquarters, Meadville, Pennsylvania.



Calvin Ernst educating the Sales Team in a field of *Oenothera fruticosa* var. *fruticosa* (Sundrops).

## Dear friends,

Fifty-six years ago, I founded my company with a small planting on my parents' farm. Ernst Conservation Seeds has since expanded to produce more than 200 species on 10,000 acres of land in northwest Pennsylvania. Our relationships with other growers and seed companies have expanded our total offerings to more than 400 native and naturalized species.

Our focus over the first 25 years was producing naturalized species for preventing soil erosion along highways and the rehabilitation of strip-mined land. Our second quarter century saw our focus pivot to producing native species for erosion control, habitat creation/restoration, and beautification. Over the last six years, we have also focused on enhancing and producing habitat for pollinators and songbirds.

I believe the coming years hold much promise but, as with all times in history, there will be challenges to address. An ongoing challenge will be how to manage crops for which no book has been written. We also face how to mechanize more of our production and maximize yields while being good stewards of the environment. For our customers, the overabundance of deer in much of the Northeast will challenge the establishment of diverse meadows that sustain a variety of wildlife. For most of our service area, invasive species will be a challenge. I am optimistic our young agronomists will develop effective control strategies for most invasive species, and I foresee additional companies forming that will be dedicated to controlling invasive species.

As we forge ahead, I see great opportunities for those involved in conservation and stewardship activities, as well as an increased interest in the use of natives for many of the purposes for which they are already being used. I anticipate we will see expanded use of buffer plantings to reduce nutrient runoff from crop fields to our rivers, lakes, and estuaries. We will continue to see increasing interest in making solar farms pollinator and songbird friendly. If our past suggests anything about our future, it is that an opportunity will arise that we are not contemplating today!

Thank you for being a part of the journey through our first 56 years. I hope we will be part of each other's lives for decades to come.

Yours in conservation,

**Calvin L. Ernst**  
President

### VISIT OUR WEBSITE

For the Wetland Indicator Status of species sold by Ernst Seeds, visit the individual species page online at [www.ernstseed.com](http://www.ernstseed.com).

**POLLINATOR-FRIENDLY SITES**  
Page 26

**WET MEADOW AND WETLAND SITES**  
Page 38

**OBSTACLES TO A SUCCESSFUL MEADOW ESTABLISHMENT**  
Page 55

**BIOMASS**  
Page 60

ERNST TEAM



**CALVIN ERNST (MARCIA ERNST)**  
*President*



**ANDY ERNST**  
*Vice President*



**MICHAEL ERNST**  
*Vice President*



**ROBIN ERNST**  
*Partner*

# THE ERNST TEAM

OUR TEAM TAKES GREAT PRIDE IN THE WORK WE DO AND THE QUALITY OF THE PRODUCTS WE SELL. FROM THE PRODUCTION FIELDS TO THE SALES OFFICE, OUR WORKFORCE IS THE BEST AT WHAT THEY DO, AND EACH MEMBER IS REGARDED AS AN EXTENSION OF THE ERNST FAMILY.



**DAN ARNETT**  
*Biomass Manager*



**STEPHANIE BRECKENRIDGE**  
*Operations and Inventory Manager*



**PAULA DITHRICH**  
*Senior Accounting Manager*



**MARK FIELY**  
*Horticulturist*



**CHERI HAINES**  
*Human Resources Manager*




**KATHY HAVEN**  
*Executive Assistant*



**NIKKI HINDLE**  
*Sales Manager*



**KEVIN JAMISON**  
*Agronomist*



**GREG KEDZIERSKI**  
*Plant Material Specialist*

ERNST TEAM



ACCOUNTING & FINANCE



PLANTING & HARVESTING



SALES & ORDER PROCESSING



BIOENGINEERING



SEED CONDITIONING



BIOMASS



MARKETING



GREENHOUSE



WAREHOUSE & SHIPPING

## DEFINING NATIVE vs. NATURALIZED SEEDS

### NATIVE

Species that existed locally prior to European settlement.

### NATURALIZED

Species not native to a certain area that grow, reproduce, and maintain themselves without interference.

### ECOTYPE

Native species found in a defined area, state, or region.

### VARIETY

A subdivision of native or naturalized species having one or more distinct, consistent, though often inconspicuous, traits.

---

## WHY DO STOCK MIX FORMULATIONS CHANGE?

Stock seed mix formulas may vary within a year or between years. Each mix is created with a particular guiding philosophy. As new species become available, they will be added to enhance the performance of the mixes.

Occasionally, a species may not be available due to crop failure or high sales, resulting in a reformulation using the remaining species in the mix. To adapt to these variations in our formulations, we recommend using the phrase "Ernst Mix (X) as currently formulated" when writing specifications.



ERNMX-140 Partially Shaded Area Roadside Mix



ERNMX-168 Northeast Annual & Perennial Wildflower Mix

# UNDERSTANDING PURE LIVE SEED (PLS)

The Pure Live Seed (PLS) standard was developed to aid in determining the appropriate amount of bulk seed to be applied. PLS refers to the percentage of live seed by weight in a seed lot having the potential to develop into a seedling. Live seed refers to the germination percent, dormant seed, and hard seed. The remaining seed in the lot contains inert matter, other crops, and/or weed seeds and is not viable.

## INDEX:

**PURE SEED:** Viable and non-viable seed of the desired species.

**% PURITY:** Percentage of pure seed by weight.

**% GERMINATION:** Percent of seed by weight having the potential to germinate relatively quickly.

**% HARD SEED:** Percent of viable seed having a hard seed coat that can take longer to break and germinate.

**% DORMANT SEED:** Percent of viable seed taking longer to germinate, other than hard seed.

**% TOTAL GERMINATION:** Percent of seed by weight composed of hard seed, dormant seed, and germination percentage having the potential to produce a seedling.

**% NONVIALE SEED:** Percent of seed that will not germinate.

**% INERT MATTER:** Material other than seed.

**VIALE SEED:** Pure seed having the potential to produce a seedling.

**WEED SEED:** Seeds or other reproductive parts of species recognized by law as weeds.

**OTHER CROP SEED:** Seeds that are not weeds or pure seed.

## HOW TO CALCULATE PLS:

**Percent Total Germination = (Germination + Hard Seed + Dormant Seed)**

**70 + 15 + 5 = 90% Total Germination**

Use that figure and the purity percentage to calculate the PLS percentage.


**Multiply Total Germination by Purity / 100 = PLS %**  
**(95% X 90%)/100 = 85.5% of Pure Live Seed**

*What Does This Mean?*

85.5% of this seed lot by weight has the potential to grow. If 10 pounds of pure live seed is needed on a site, the quantity of bulk seed needed is determined by:

**100 / PLS = Pounds of bulk seed needed to produce 1 lb of PLS**

**100 / 85.5 = 1.16 lbs of bulk seed X 10 lbs = 11.6 lb of bulk seed**



# EXPECTATIONS OF NATIVE SPECIES



# EXPECTATIONS OF NATIVE SPECIES

GERMINATION AND GROWTH (all the following assume adequate light, adequate soil temperature, appropriate moisture, and good seed-to-soil contact):

In general, annual species have less seed dormancy than biennials and biennials have less than perennials. Seed dormancy is nature's hedge against unfavorable conditions during a plant's life cycle. Unfavorable conditions can be, but are not limited to, late spring frost or drought. Dormant seeds germinate when favorable conditions are present.

## ANNUALS

Most species germinate, flower, and set seed by the end of the first full growing season. Germination of an individual species is likely to be high.

## BIENNIALS

Most species germinate, with some plants within a species' population flowering and setting seed in the first full growing season. The bulk of the plants will flower and set seed in the second growing season. Germination of an individual species is likely to be lower than that of an annual due to seed dormancy.

## PERENNIALS

**1. Warm Season Grasses:** Germination will occur in the spring when moisture conditions are appropriate and soil temperatures exceed 55°F (12°C) at a 3" depth. Best germination occurs when soil temperatures are much higher.

Most of these species do not require cold, wet stratification to produce an adequate stand. Two exceptions are *Tripsacum dactyloides* (Eastern Gamagrass), which requires 14-60 days of stratification, and *Chasmanthium latifolium* (River Oats), which requires 60 days of stratification for northern genotypes. Stratification is the process by which seed is exposed to cool, moist conditions.

While cold, wet stratification is not necessary in most cases to produce an adequate stand, 20%-50% of the seed may be dormant. Most seedlings that emerge will be growing by the end of the second full growing season.

Greatest growth of these species occurs when air temperatures are 75°F-95°F (24°C-35°C). Most of the growth is in root development the first season. Very few (<5%) plants within a species may flower and set seed in the first growing season. Maximum plant development may take two years or longer.

**2. Cool Season Grasses:** Some species will germinate when temperatures are a little higher than 40°F (4°C) while others will require warmer temperatures. They may germinate in the fall or spring. Adequate stands of most species do not require stratification; however, 50% of the seed may remain dormant without stratification. Most seedlings that emerge will be growing by the end of the second full growing season.

Greatest growth occurs when temperatures are 65°F-85°F (18°C-29°C). With adequate moisture and nutrients, some flowering and seed set may occur in the first growing season.



New growth of *Tripsacum dactyloides* (Eastern Gamagrass) in a Florida field in the spring.



*Tripsacum dactyloides* (Eastern Gamagrass).

## EXPECTATIONS OF NATIVE SPECIES



**3. Some sedges** (*Carex albolutescens*, *scoparia*, *vulpi-noidea*), **rushes** (*Juncus effusus*, *tenuis*) and **bulrushes** (*Scirpus atrovirens*, *cyperinus*, *expansus*, *polyphyllus*) have a very high seed count per pound of seed. When planted in the spring, a substantial number of seedlings may be produced by these species in the first growing season. These seedlings may represent 5% or less of the total seeds present. Flowering and seed production will occur one to two growing seasons after an individual seedling has germinated. Maximum germination will take at least two years due to seed dormancy. Sedges and bulrushes are recognizable by the arrangement of any three successive leaves in a pattern resembling the spokes in the Mercedes™ symbol. *Juncus spp.* will have round stems that originate at a common point near or on top of the soil.

**4. Some sedges** (*Carex comosa*, *crinita*, *folliculata*, *frankii*, *glaucescens*, *grayi*, *granularis*, *intumescens*, *lupulina*, *lurida*, *squarrosa*, *stipata*, *stricta*) and **bulrushes** (*Scirpus acutus*, *americanus*, *robustus*, *validus*) have a high level of seed dormancy and may not have consequential germination without stratification.

Most seedlings will emerge in the first and second growing seasons after stratification (artificially or naturally). Plants will flower and set seed one to three years after they germinate. *Carex spp.* in this group may be recognized as described above for other *Carex spp.* *Scirpus spp.* in this group have round or triangular stems arising from a point often below the soil surface. The stems are typically larger than those of *Juncus spp.*



**5. For most broadleaf species**, some germination will occur in the first year without stratification (artificial or natural). A high percentage of species and seeds within the species are likely to germinate in the first growing season following the first winter *in situ* (on-site). Most of the seeds that germinate will have done so by the end of the growing season following stratification. Following germination, blooms may occur in the first growing season: *Heliopsis helianthoides* (Oxeye Sunflower); second growing season: *Rudbeckia triloba* (Browneyed Susan), *Aster spp.*, *Monarda spp.*, *Penstemon spp.*, *Solidago spp.*; after three to five growing seasons: *Liatris spp.*; or, not until the seventh growing season: *Baptisia tinctoria* (Yellow False Indigo). The number of blooming years depends on soil fertility, available moisture, and growing season temperatures. It may be shorter for a given species the further south one is located.

## EXPECTATIONS OF NATIVE SPECIES



**6. Seed dormancy in perennial species** is affected by latitude of ecotype origin. In greenhouse studies, we have found that northern ecotypes (PA, OH, NY, NJ) typically require more weeks of cold, wet stratification than southern ecotypes (FL, GA, NC, SC) of the same species.

Most of our native seed mixes are composed of perennial species. Mixes dominated by perennial species have the potential to last more than a decade if properly maintained. For all mixes, a site must be kept free from invasive species or aggressive weeds. Mixes of herbaceous species with no tree, shrub, or vine components in the formula must be kept free from the encroachment of woody or vine species with controlled burning, mowing, or spot spraying.

The natural communities we create with native seed mixes are dynamic. Annuals, biennials, and short-lived perennials may be widely present in the landscape in the first three growing seasons, but non-existent or present in small pockets by the fifth growing season. Over time, colonies of some long-lived perennials

will grow larger in area and species composition will change in response to annual rainfall variations.

It is not unusual for those new to planting meadows to be nervous about a mix's performance during its establishment year. Typically, customers need confirmation that the desirable species are growing. Fortunately, our ability to assess a situation is assisted by a small set of species that generally germinate very well.

For wetland meadows, some common early emerging species include: *Asclepias incarnata* (Swamp Milkweed), *Eupatorium perfoliatum* (Boneset) and *Carex spp.* For upland meadows, some common early emerging species include: *Chamaecrista fasciculata* (Partridge Pea), *Elymus virginicus* (Virginia Wildrye), *Helianthus angustifolius* (Narrowleaf Sunflower), *Heliopsis helianthoides* (Oxeye Sunflower), *Monarda fistulosa* (Wild Bergamot), *Penstemon digitalis* (Tall White Beardtongue), and *Rudbeckia hirta* (Blackeyed Susan). Seedling images of many of these species are available at the download center on our website.

---


### DISCLAIMER

The information in this review of practices is the result of over 50 years' experience in seed production. Ernst Conservation Seeds has been supplying seeds and consulting in the reseeded of tens of thousands of acres of roadsides, surface-mined lands, conservation, and restoration sites in eastern North America, as well as growing and supplying seed and consulting in the planting of hundreds of thousands of acres of CRP/CREP-related areas for erosion control and wildlife habitat.

All these practices are opinion only and our best advice as a result of these experiences. These recommendations are for individual consideration and do not cover all the conditions that will be encountered in the field.

Ernst Conservation Seeds is not responsible for conditions that will be encountered in individual situations. The use of brand names does not represent our endorsement of a specific product; rather, it represents our experience only and has not necessarily been replicated in peer-reviewed research. The use of chemical pest control agents is subject to manufacturers' instructions and labeling, as well as federal, state, and local regulations.

# ESTABLISHMENT GUIDE





A pasture buffer in Meadville, Pennsylvania.

**ESTABLISHMENT GUIDE INTRODUCTION**

In eastern North America, there is a wide variety of native vegetation to replicate. Most planting objectives fall into the following categories:

- › Soil erosion control & soil stabilization on slopes and along waterways
- › Beautification & enhancement of landscapes
- › Biodiversity & wildlife habitat enhancement and restoration
- › Bioremediation to correct environmental disturbances
- › Historical, cultural & ecological restoration
- › Habitat for honeybees & native pollinators (butterflies, bumblebees, etc.)
- › Native species for renewable biomass production

Using native plants saves time and money while improving ecological function. Reduced water, chemical, fertilizer, and maintenance needs make them a sustainable and environmentally sound choice for virtually all scenarios. Select a mix of species that creates the desired outcome for the project. Goals should be compatible with site conditions that cannot be altered. Native plant communities can be selected to meet nearly all site conditions.

Please review the appropriate section(s) below for information regarding seed mix selection and seeding methods. Matching the functional goals of the site and

site conditions to the appropriate seed mix will lead to greater project success. The stock seed mixes noted in each section represent a mere sampling of our complete list of mixes. A more comprehensive list may be found at [www.ernstseed.com](http://www.ernstseed.com) or by contacting a member of our sales team. Mixes can also be customized to specific needs as well as those of a site and ecological region.

Upland & Meadow Sites	<b>19</b>
Wildlife Habitat & Food Plot Sites	<b>23</b>
Pollinator-Friendly Sites	<b>26</b>
Riparian Sites	<b>29</b>
Disturbed Sites & Steep Slopes	<b>32</b>
Utility Right-of-Way Sites	<b>36</b>
Wet Meadow & Wetland Sites	<b>38</b>
Partially Shaded Sites	<b>41</b>
Stormwater Management Sites	<b>44</b>
Southeastern U.S. Sites	<b>48</b>
Biomass Production Sites	<b>52</b>



Loading one of the seed hoppers on a Truax no-till drill during spring planting.

## FALL VS. SPRING SEEDING

Traditionally, seeding is thought of as a spring activity, but it can also occur during the dormant season. Fall seeding works well for restoration projects completed in the summer. While there are some noteworthy advantages to fall seeding, seeding in either spring or fall will produce good results. In drought-prone regions, seeding should be timed to take advantage of the available moisture in the area.

### FALL OR “DORMANT” SEEDING

- › Fall seeding imitates natural reseeding. Dormant seeding can take place when soils are dry enough to work.
- › Good seed-to-soil contact occurs through precipitation and the freeze-thaw cycle.
- › Natural stratification and scarification occur; natural changes within the seed or to the seed coat during the winter enhance germination in the spring.
- › Mulching is an important element of dormant seeding to protect the soil.
- › Some seed may be lost to decay and wildlife consumption during the winter.
- › Establishment may be hindered by growth of winter annuals in the fall.

### FROST SEEDING

- › Frost seeding is the act of broadcasting seed onto or drilling into frozen soil.
- › Good seed-to-soil contact occurs through the freeze-thaw cycle.
- › Natural stratification and scarification occur; natural changes within the seed or to the seed coat during the winter enhance germination in the spring.

- › Mulching may be done.
- › Some seed may be lost to decay and wildlife consumption during the winter.
- › Establishment may be hindered by growth of winter annuals in the fall.

### SPRING SEEDING

- › Cool season species germinate soon after seeding.
- › Germination of warm season species generally occurs within three weeks of the soil temperature reaching 55°F (13°C).
- › Seed loss due to decay and wildlife consumption is minimized.
- › Seed-to-soil contact should be accomplished by working the seed into the soil 1/4”-1/2” deep.
- › Seeding may be delayed until weed control is applied to improve establishment.
- › Irrigation during dry weather periods is necessary for proper germination.
- › Light mulching is an important element of seeding to protect both the seed and soil and retain moisture.
- › When planting native grasses for biomass, seeding generally takes place during the spring when soil temperatures at a 3” depth are near 55°F and rising.

## SEEDING METHODS



### DRILL SEEDING

Drill seeding is a mechanical means of creating furrows (openings) in the soil and metering seed in at a uniform rate. A drill seeder is practical for seeding multiple acres in larger areas.

Conventional drills can work in tilled and partly tilled soil. No-till drills are designed to work in soil that has not been tilled. They have heavy openers that cut through vegetation and sod to make a furrow for seed placement. With the proper adjustment, a no-till drill can work in tilled soil. It has discs that aid in loosening the soil. All drills should be equipped with a closing or packing wheel that follows seed placement.

The goal of drill seeding is to achieve uniform seed distribution over the site with seed placement at the correct depth (1/4"-1/2") and good seed-to-soil contact. Calibrating a drill or broadcast seeder depends on seed bulk density and required application rates. Manufacturers provide manuals with charts to guide seeding rate calibration. To ensure uniform application of seed, conduct a test run over a small area using the appropriate amount of seed for that area, then make any necessary adjustments.

Most traditional seed drills are designed to handle seeds with high bulk densities, such as oats and wheat. Some drills may have a small seed box able to plant small seeds, such as alfalfa, clover, and switchgrass.

Many native and naturalized species are fluffy and will not readily flow through a traditional seed drill. Examples of fluffy seed include little bluestem, big bluestem, and indiagrass. With the aid of a bulking agent, some fluffy seeds may be planted through the large seed box of a traditional drill. Bulking agents include kitty litter, dry sawdust, vermiculite, or rice hulls. Test with a small amount of seed. Native seed drills, such as Truax, have specialized seed boxes that are effective for planting fluffy seed. When seed will not readily flow through a native seed drill's fluffy seed box, a bulking agent may be needed.



### HAND SEEDING

Hand seeding is the casting of seed onto the soil. Hand seeding is used on small plots or difficult terrain where seeding with machinery is not an option. The goal is to achieve an even distribution of seed over the site. This can be accomplished by spreading half of the seed in one pass and the balance in a perpendicular pass. To ensure uniform application of seed, conduct a test run over a small area using the appropriate amount of seed for that area. To know how wide to make your passes, check the width of seed distribution. If possible, a light raking to a depth of 1/4" and/or firming with a lawn or Brillion-type roller is recommended to achieve good seed-to-soil contact. Cover with straw mulch at 70 lb per 1,000 sq ft or hydromulch at 34 lb per 1,000 sq ft. When the volume of seed to be applied is small (less than 50 lb per acre), a bulking agent may be helpful to provide the volume necessary to get uniform application. Such bulking agents include kitty litter, dry sawdust, vermiculite, or rice hulls.

## SEEDING METHODS



### BROADCAST SEEDING

A broadcast seeder consists of a hopper with an adjustable door that regulates seed flow onto a spinner. Some broadcast seeders have an agitator that aids with seed flow in the hopper. Broadcast seeders are commonly used to spread seed, fertilizer, lime, and other granular products. The goal is to achieve an even distribution of seed over the site. To ensure uniform application of seed, conduct a test run over a small area using the appropriate amount of seed for that area. To know how wide to make the passes, check the width of seed distribution from the spreader. The settings can then be adjusted as needed. To achieve better distribution, spread half of the seed in one pass and the balance in a perpendicular pass. We recommend refilling the hopper when it is 1/3 full rather than letting it empty out. Follow up by tracking or firming the seed into the soil with a lawn or Brillion-type roller to achieve good seed-to-soil contact. Do not roll or track the seed if the soil is wet. Cover with straw mulch at 70 lb per 1,000 sq ft or hydromulch at 34 lb per 1,000 sq ft. Many native seeds are fluffy and will not flow uniformly through a broadcast seeder. To enhance the flow, mix the seed with a bulking agent of similar density. Dry sawdust, vermiculite, or rice hulls are some options. An agitator in the hopper may be required in these circumstances. We recommend a minimum rate of 50 lb per acre of seed and bulking agent. A bulking agent can also be helpful if you are planting small quantities of seed. It provides the volume necessary to get uniform application. For fine seeds, kitty litter is a more appropriate bulking agent.

### CULTIPACKING

A cultipacker is an excellent way of covering the seed with a minimum amount of soil to ensure proper seed-to-soil contact. It resembles a large rolling pin with evenly spaced ridges and dimples. The primary functions of a cultipacker are to break up clods, remove excess air spaces from loose soil, and smooth the soil. The heavy-duty smooth, spoke, or crowfoot rollers provide clod-breaking and smoothing capabilities. As with any tillage, it is important not to overwork the soil or work it when it is too wet.



### STRAW MULCHING

A straw-mulch blower distributes mulch over a seeded area. It has a slide (or chute) in which to feed the mulch, chopper blades to break up the mulch, and a blower to spread the mulch over large areas. Straw mulch may be spread by hand in smaller areas. It is important to use weed-free straw from small grains, such as oats or grain rye, to minimize potential weed issues.



### HYDROSEEDING

A hydroseeder combines water, seed, fertilizer and, sometimes, hydromulch into a mix that is pumped through a nozzle and sprayed uniformly over the area to be seeded. Hydroseeders can distribute this mix at 150' or more, allowing for the ability to seed terrain that may not be accessible with other seeding methods, such as steep slopes, roadside cuts, or sites that are too wet. Using hydromulch aids in seed placement and reduces erosion on slopes. Depending on site conditions, use of erosion control blankets or straw mulch may be needed to cover the seed. Many native seeds should be broadcast with 500 lb per acre of mulch as a marker. Do not exceed this amount as native seeds may die if suspended in the mulch with little or no seed-to-soil contact. The balance of the hydromulch, often 1,000 lb per acre, may be applied on top in a secondary application.

## TOOLS FOR PREPARATION



### MINIMUM-TILL EQUIPMENT

Minimum-till equipment incorporates a portion of the surface vegetation into the soil and levels uneven surfaces. One of the most common tools is a disc which cuts through vegetation, sod, or hard soil and partially turns or tills it into the soil. Similar equipment that turns part of the vegetative residue into the soil is known as Aerway® or Turbo® Till.



### CHISEL PLOW

A chisel plow is primarily used to break up hardpan soil or loosen compacted soil while leaving a high percentage of debris on top. It is a minimum-till plow because it does not dislodge or turn over the entire soil profile the way a moldboard plow does. A chisel plow may be adjusted to till shallow or deep and typically has C-shaped shanks mounted on dual coil springs. The frame, shanks, and springs are of sufficient weight, size, and strength to provide a cutting depth of 8"-12". To make the soil smooth enough for planting after the use of a chisel plow, use a disc harrow, tandem disc harrow, or offset disc harrow of sufficient weight and size to provide a cutting depth of 6"-8".



### ROTOTILLER

A rototiller pulverizes the soil with rotating blades and incorporates soil amendments and surface vegetation. Most units till up to 6" deep.



### TRACKING

Tracking is the use of a crawler or rubber-tired tractor to make depressions and firm loose soil after construction or tilling. Tracks should be oriented perpendicular to the slope of a site. Tracking depressions aid in reducing erosion and retaining seed and moisture. The firm, but not compacted, seedbed will not dry out as quickly as loose soil.

## TOOLS FOR MAINTENANCE



### ROTARY MOWER

A heavy-duty rotary mower can be utilized as a brush hog to tame heavy grass and light brush, such as multiflora rose, honeysuckle, and small tree seedlings, on under-utilized fields difficult to mow with a discbine or sickle bar mower.



### DISCBINE MOWER

A discbine mower is a hay-harvesting machine with high-speed rotary discs for mowing and baling biomass and assembling the material into a windrow.

## WHEN TO MOW

Following the establishment year, typically, mowing during the growing season should not be necessary unless it is in lieu of herbicides to control weeds. Mowing height should be no lower than 8”.

To prevent succession of woody species in an established meadow, an important aspect of a maintenance program is an early spring mowing close to the ground (2”). Mowing should occur every one to three years in late winter or early spring and shortly before spring nesting season. Spring mowing will leave food and cover for wildlife through the winter without disrupting nesting of grassland birds.



### SPRAYER

Sprayers come in various sizes and styles, including common hand-held units like the one shown here. They are often preferred for carefully targeted spraying of unwanted or invasive vegetation. Larger areas may be sprayed effectively using tractor or ATV-drawn tank units.

The use of herbicides for controlling undesirable vegetation can be an important part of an integrated pest management (IPM) program when applied according to the manufacturer’s label. Prior to using any herbicide, read the label for safe handling and application information. Many herbicides are only available to licensed applicators. In these cases, a licensed professional should be employed.

# UPLAND & MEADOW SITES



**UPLAND SITES ARE** characterized as being dry most of the year. Soils on these sites are well-drained and may consist of sandy clay, sandy loam, loam, or shale. The topsoil layer may be thin and subject to drought. If the site is sandy, refer to the Southeastern U.S. Sites Establishment Guide, p. 48. Upland examples: Naturally rocky soil subject to erosion or steep road cuts. Meadow examples: Abandoned farm fields, lawns, vacant land, or roadsides.



An upland meadow using ERNMX-179 Butterfly & Hummingbird Garden Mix. Credit: Paul Stead



**HABITAT:**  
Upland to mesic sites, typically in full sun for at least half the day with good air circulation.



**FERTILITY:**  
Natural fertility is usually adequate; therefore, fertilizer and lime are not needed (fertilizer often aids in the growth of weeds and invasives). Check soil pH and select species adapted to that pH.



**SEEDING METHOD:**  
Hand seed, broadcast seed, hydroseed, or drill seed.

### SITE PREPARATION

If the site was previously a lawn or crop field to which herbicides were applied, it is important to allow the appropriate time interval for the herbicide residues to break down prior to planting. Some herbicide residues can prevent seedlings from germinating.

Competition from invasive or undesirable vegetation is the most limiting factor for a successful upland meadow establishment. Prior to planting, all such vegetation must be fully controlled. Typical control strategies include repeated tilling, smothering with black plastic, or herbicides. While any of these methods may control existing weeds, they will not kill all weed seeds lying dormant in the soil. Seeds of such species as velvetleaf and pokeweed may germinate many decades after the species last flowered on the site.

When using the tillage strategy, a site is disc harrowed every two to four weeks for one to two months. The underlying principle of this process is that the root system of perennial species will be worn out to the point of killing the species. In addition, tillage will stimulate germination of some dormant weed seeds that will be killed with subsequent tillage. Planting should not take place until perennial species are completely killed.

Black plastic may also be used to kill weeds. It may be laid across tilled or untilled soil and anchored down by burying the edges in soil or laying boards or bricks across the surface. This strategy should be utilized in a growing season when the intent is to fall plant in the same year or spring plant the following year.

Use of an approved herbicide, such as glyphosate (Roundup® or Rodeo®), by a licensed spray technician is the most common and least time-intensive method for controlling existing vegetation. Since herbicides are most effective on actively growing plant tissues, they are very effective on new spring growth. Spraying should begin when growth reaches 6". A follow-up application one to two weeks later will address skips or persistent species. If substantial plant tissue remains on the surface following a full kill by herbicides, close mowing, tillage, or burning may be necessary to achieve good seed-to-soil contact.

## GROWING SEASON MAINTENANCE

### MAINTENANCE

- › Problem weeds should be hand pulled in annual wildflower mixes or annual & perennial wildflower mixes. For all other mixes, see below.

### FIRST GROWING SEASON

- › Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a brush hog mower or string trimmer. Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed. Trimming should cease by mid-September.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Stinger®, or Milestone®. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

### SECOND & SUBSEQUENT GROWING SEASONS

- › Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any remaining material from the previous year close to the ground (approximately 2"). This will allow the soil to warm more quickly, stimulating emergence and growth of native seedlings and reducing the likelihood of shrub invasion.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Stinger®, or Milestone®. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

### SPECIAL CIRCUMSTANCES - SECOND GROWING SEASON

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". Trimming should cease by mid-September. To prevent weed reinfestation, use of an appropriate selective herbicide in conjunction with a seed mix tolerant of that herbicide may be necessary. \*



A wildflower meadow using ERNMX-168 Northeast Annual & Perennial Wildflower Mix. Credit: Joseph Garmon

UPLAND & MEADOW SITES



A meadow installation at a Princeton, New Jersey residence. Credit: OPEN Landscape Architecture

UPLAND & MEADOW SEED MIXES	
ERNMX-105	Northeastern U.S. Roadside Native Mix
ERNMX-110	Ernst Native Biomass Mix for Strip Mines & Natural Gas Production Sites
ERNMX-111	Ernst Native Habitat for Strip Mines Mix
ERNMX-115	Biodiverse Polyculture Mix for Biomass Production & Wildlife Habitat
ERNMX-117	Warm Season Grass Mix
ERNMX-123	Native Upland Wildlife Forage & Cover Meadow Mix
ERNMX-125	Northeastern U.S. Roadside Native Mix without Grasses
ERNMX-153	Showy Northeast Native Wildflower & Grass Mix
ERNMX-153-1	Showy Northeast Native Wildflower Mix
ERNMX-156	Low-Growing Wildflower & Grass Mix
ERNMX-166	Plateau-Tolerant Wildflower & Grass Mix
ERNMX-166-1	Plateau-Tolerant Wildflower Mix
ERNMX-167	Annual Wildflower Mix
ERNMX-168	Northeast Annual & Perennial Wildflower Mix
ERNMX-169	Southeast Annual & Perennial Wildflower Mix
ERNMX-170	Annual & Perennial Wildlife Food Plot Mix
ERNMX-171	Multi-Purpose/Multi-Year Wildlife Food & Shelter Mix
ERNMX-172	Maryland Upland Mix
ERNMX-173	Eastern Native Habitat & CREP Mix
ERNMX-174	Virginia Gentleman's Mix
ERNMX-177	Eastern Ecotype Native Grass Mix
ERNMX-179	Butterfly & Hummingbird Garden Mix

**THESE MIXES ARE USED IN FULL SUN WITH WELL-DRAINED SOILS AND PROVIDE FOOD AND/OR COVER FOR WILDLIFE. MEADOW AND WILDFLOWER MIXES PROVIDE FOOD FOR INSECTS, INCLUDING NATIVE POLLINATORS.**

**VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.**

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See "Disclaimer," p. 11. For "Expectations of Native Species," see p. 8.

# WILDLIFE HABITAT & FOOD PLOT SITES



A ringneck pheasant rooster emerges from a grassy woodland edge.

**WILDLIFE FOOD PLOT SITES ARE GENERALLY** small clearings in wooded areas or farmland. While planting dates differ between annual and perennial mixes, most are designed for spring or fall planting. Examples: Woodland openings, remote areas of large residential lots, abandoned fields, or timber harvest and loading areas.



A young whitetail deer in *Andropogon virginicus* (Broomsedge).



**HABITAT:**

Well-drained or moderately well-drained sites in woodland openings (ideally near water sources); typically in full sun for at least half the day; upland species may be planted.



**FERTILITY:**

Check soil pH and fertility; adding lime can improve the nutritional value of vegetation beneficial for wildlife. If a soil test has not been obtained, a starter fertilizer, such as 200 lb per acre of 16-16-16, should be applied. If uncertain about the soil pH, add 1,000-2,000 lb of lime per acre which will provide plants with essential nutrients without pushing up a lot of top growth. Lime and fertilizer may be incorporated into the soil using a tiller. After incorporating amendments, smooth the soil to develop a good seedbed. If broadcasting the seed, run a spring-tooth harrow or ATV over the site to incorporate the seed into the soil 1/4"-1/2" deep.



**SEEDING METHOD:**

Hand seed, broadcast seed, or drill seed.

**SITE PREPARATION**

If the site was previously a lawn or crop field to which herbicides were applied, it is important to allow the appropriate interval for the herbicide residues to break down prior to planting. Some herbicide residues can prevent seedling germination.

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide, such as glyphosate (Roundup® or Rodeo®), or by tilling the weeds into the soil. Hand pull or spot spray problem weeds. Perennial weeds not addressed before establishment will be difficult to remove later. Good pre-seeding weed control may require repeated tilling or spraying two applications of glyphosate at least two weeks apart.

**GROWING SEASON MAINTENANCE**

**GENERAL MAINTENANCE**

› It may be necessary to mow some mixes to 4"-6" in order to keep plants young and tender. Annual food plot mixes must be planted yearly. 🌱



ERNMX-184 Fall Sweets Wildlife Mix. Credit: Elicia Winner



A ringneck pheasant rooster takes flight in a hunting preserve using Ernst habitat mixes.



A pair of eastern wild turkey jakes in a spring food plot.

## WILDLIFE HABITAT & FOOD PLOT SEED MIXES

ERNMX-130	Wildlife Food Plot Mix
ERNMX-133	Keystone Deer & Turkey Habitat Mix
ERNMX-133-1	Keystone Big Buck Mix
ERNMX-170	Annual & Perennial Wildlife Food Plot Mix
ERNMX-171	Multi-Purpose/Multi-Year Wildlife Food & Shelter Mix
ERNMX-184	Fall Sweets Wildlife Mix
ERMNX-185	Spring Greens Mix

THESE MIXES PROVIDE FOOD FOR DEER, TURKEY, AND OTHER UPLAND GAME BIRDS.

VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See “Disclaimer,” p. 11. For “Expectations of Native Species,” see p. 8.

# POLLINATOR-FRIENDLY SITES



Carpenter Bee (*Xylocopa virginica*) enjoying the fall bloom of *Solidago speciosa* (Showy Goldenrod).

**POLLINATOR MEADOWS MAY BE USED TO** attract honeybees and more than 4,000 species of native pollinators in North America, including bees, butterflies, hummingbirds, and even some flies. Increased attention to the plight of such pollinators as honeybees and monarch butterflies by the government and private sector has spurred an interest in developing pollinator-friendly habitats across the North American landscape. One of the highest examples was the 2015 White House announcement of the National Strategy to Promote the Health of Honeybees and Other Pollinators.



Hummingbird Clearwing (*Hemaris thysbe*) visiting *Monarda fistulosa* (Wild Bergamot) at the Erie National Wildlife Reserve in Guys Mills, Pennsylvania.

## ABOUT POLLINATOR-FRIENDLY SITES

Almost any site not intended to be mowed repeatedly during the growing season may be designed to be pollinator friendly. Pollinator-friendly meadows are often thought of as being planted on flat sites in full sun, but they can also be planted at woodland edges. Pollinator-friendly species aid in soil stabilization on steep slopes and riparian areas. Storm basins and wetlands are more aesthetically pleasing when enhanced with pollinator-friendly species. Rights-of-way under utility transmission lines, above pipelines, and along roadways can be developed to an ecologically beneficial state when functional diversity and pollinator-friendly species are incorporated into seed mix design. When planted within a solar array, transpiration from native plants can reduce panel temperatures thereby increasing panel efficiency.

The primary energy source for most adult bees, butterflies, and other flower-loving pollinators is nectar. Pollen is essential for providing proteins and lipids to developing bee larvae while leaf tissue from specific host plant families is required for butterfly caterpillars. Most native bees are nectar generalists in that, though pollen specialists, they can consume nectar from many plant families. Pollen that the larvae will consume requires a specific ratio of proteins to lipids. The best sources of pollen for native bees as well as leaf tissue for native



Solar power generation sites can be sources for pollinator habitat.

butterflies are the native plant species with which they have co-evolved.

While not native to the U.S., honeybees have evolved to be able to use pollen from a wide range of species. Like native bees, honeybees feed nectar and pollen to their larvae. They also need pollen to have a particular protein-to-lipid ratio that they get by collecting pollen from a variety of plant species.

Adult pollinators do not have the same dietary needs as when in the juvenile stage. To meet the dietary needs of a wide range of pollinators, it is important to know



Tachinid Fly on *Pycnanthemum incanum* (Hoary Mountainmint).

that some pollinator species are not active for the entire growing season. During the active period, food and nesting resources must be available. The availability of flowering shrubs or trees for pollen and/or nectar before herbaceous species bloom in the spring is beneficial to some pollinator species. Continuity of bloom from as early in the season to as late in the season as possible is important. A minimum of three species should be in bloom in the spring, summer, and fall, with a total of 26 blooming species being ideal. These species should represent five or more plant families. For the benefit of monarchs, milkweeds should be planted.

While dietary needs of honeybees have much in common with that of native bees, an important difference is that honeybees are attracted to masses of bloom. For that reason, rather than planting a mix of 26 species, multiple mixes should be planted, each with three to four blooming species with no overlapping of bloom within a mix. Bloom overlap will be provided by the other mixes. \*

**TO SUPPORT THE GREATEST DIVERSITY OF NATIVE POLLINATORS:**

- > Provide continuity of bloom from as early to as late in the season as possible.
- > Minimum of three species blooming in spring, summer, and fall.
- > Twenty-six blooming species from 5 or more plant families.
- > Plant milkweeds for monarchs.

POLLINATOR-FRIENDLY SITES SEED MIXES	
ERNMX-105	Mesic to Dry Native Pollinator Mix
ERNMX-125	Mesic to Dry Native Pollinator Mix without Grasses
ERNMX-153	Showy Northeast Native Wildflower & Grass Mix
ERNMX-153-1	Showy Northeast Native Wildflower Mix
ERNMX-155	Deer-Resistant Meadow Mix
ERNMX-157	Honeybee Forage Mix
ERNMX-179	Butterfly & Hummingbird Garden Mix

THESE MIXES PROVIDE FOOD FOR NATIVE POLLINATORS, AS WELL AS FOOD AND COVER FOR SONGBIRDS AND GROUND NESTING BIRDS, SUCH AS TURKEY AND QUAIL.

VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See "Disclaimer," p. 11. For "Expectations of Native Species," see p. 8.

# RIPARIAN SITES




A riparian site with a streambank restoration planted mainly in *Salix* (Willow) species.

**RIPARIAN SITES ARE USUALLY** adjacent to rivers and waterways with soils often containing clay, high amounts of organic matter, and/or saturated sand. Examples: River and streambanks or damp floodplains of rivers and streams.



BEFORE



Bowman's Creek re-setting and rehabilitation project in northeastern Pennsylvania. Credit: Adam Nordfors

AFTER



**HABITAT:**

Varies from partial shade to full sun; subject to flooding; generally populated with riparian and floodplain species.



**FERTILITY:**

Due to the potential for water contamination, lime and fertilizer are not recommended. When topsoil has been depleted or removed, we recommend the addition of organic matter (straw, compost, mulch, leaf litter, etc.). Check soil pH and select species adapted to that pH.



**SEEDING METHOD:**

Hand seed, broadcast seed, or hydroseed.

**SITE PREPARATION**

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide, such as glyphosate (Rodeo®), triclopyr (Garlon® 3A), or a similar aquatic herbicide formulation, to control such undesirable vegetation as multiflora rose, honeysuckle, and woody species. CAUTION: Some persistent species, such as purple loosestrife, phragmites, Japanese knotweed, or reed canarygrass, may require multiple applications of glyphosate or triclopyr. Perennial weeds not addressed before establishment will be difficult to remove later. Before seeding, excess dead vegetation should be removed, turned under, or burned if conditions permit. Newly constructed riparian sites should be seeded as soon after construction as possible.

**GROWING SEASON MAINTENANCE**

**FIRST GROWING SEASON**

- › Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a brush hog mower or string trimmer. Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.
- › If bioengineering materials were used on the site, mowing should be above the new growth of these materials. Trimming should cease by mid-September.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, or Garlon®3A. Be vigilant in controlling invasive vines, such as bindweed, mile-a-minute, and Japanese hops. These are more easily pulled early than after two to three months of growth. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.



ERNMX-178 Riparian Buffer Mix along a small stream corridor.

**SECOND & SUBSEQUENT GROWING SEASONS**

- › Prior to new spring growth reaching 2” (e.g., shortly after forsythia or redbud blooms), trim any remaining material from the previous year close to the ground (approximately 2”). This will allow the soil to warm more quickly, stimulating emergence and growth of native seedlings and reducing the likelihood of shrub invasion.
- › If bioengineering materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Rodeo® or Garlon® 3A. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.

**SPECIAL CIRCUMSTANCES - SECOND GROWING SEASON**

- › If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8”. Trimming should cease by mid-September.

**GENERAL MAINTENANCE**

- › Grassy weeds or persistent perennials can re-establish in these soils. Monitor and control weeds by hand pulling or spot spraying. ☼

RIPARIAN SITES SEED MIXES	
ERNMX-154	Floodplain Mix
ERNMX-178	Riparian Buffer Mix

THESE MIXES ARE GOOD FOR VEGETATING RIPARIAN CORRIDORS OR FLOODPLAINS AND PROVIDE FOOD AND/OR COVER FOR INSECTS, BIRDS, AND ANIMALS. VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.


See “Disclaimer,” p. 11. For “Expectations of Native Species,” see p. 8.

# DISTURBED SITES & STEEP SLOPES



A steep slope and retaining wall utilizing ERNMX-181 Native Steep Slope Mix with Annual Ryegrass at the Millcreek Mall in Lancaster, Pennsylvania.

**DISTURBED SITES & STEEP SLOPES HAVE** various soil types and conditions typically distinguished by lower quality soils and a predisposition to runoff and erosion. Examples: Landfills, surface mines, road cuts, or construction sites.



ERNMX-181 Native Steep Slope mix with Annual Ryegrass in Morgantown, West Virginia.



**HABITAT:**

Various soils with exposed clay, sand, and rock outcropping without topsoil as a result of construction; generally populated with upland species.



**FERTILITY:**

Typically low in fertility; therefore, adding topsoil or organic matter (compost) can be very beneficial. Check soil pH and select species adapted to that pH. Add lime and fertilizer as recommended by soil analysis. Incorporate amendments into the soil in a way that will leave the soil rough and minimize soil erosion and rapid runoff (e.g., tracking). If there is a weed problem, fertilizing is not recommended.



**SEEDING METHOD:**

Hand seed, broadcast seed, hydroseed, or drill seed. For areas with slope less than 3:1, cover the seed 1/8"-1/4" deep by dragging with a spring-tooth harrow or firmly pressing the seed into the soil using a cultipacker, lawn roller, or ATV.

**SITE PREPARATION**

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide. Perennial weeds not addressed before establishment will be difficult to remove later. Whenever possible, regrade the site to reduce slope and build diversions to reduce erosion and minimize seed loss.

For areas with slope greater than 3:1, final tracking should be perpendicular to the slope. The tracks will aid in reducing erosion and retaining seed and moisture.

Mulching with straw, hydromulch, or straw/coconut fiber mats is recommended on these sites to protect the seed from drying out or washing away. For areas steeper than 3:1, the use of erosion control blankets or flexible growth medium (e.g., Flexterra®) is recommended. When using erosion control blankets, be sure they are toed in at the top of the slope.

**GROWING SEASON MAINTENANCE**

**FIRST GROWING SEASON**

› Post-planting maintenance will provide improved results if the ground is not too rough or steep. Whenever canopy height (overall vegetation) reaches 18"-24", use a brush hog mower or string trimmer to



trim the meadow to 8". Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.

- › If bioengineering materials were used on the site, mowing should be above the new growth of these materials. Trimming should cease by mid-September.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Stinger®, or Milestone®. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

## DISTURBED SITES & STEEP SLOPES SEED MIXES

ERNMX-101	Non-Native Ernst Best Strip Mine & Gas Production Mix
ERNMX-102-1	Pipeline Mix with Switchgrass
ERNMX-103	Non-Native Good Value Mine Mix
ERNMX-104	Quick Erosion Control Cover Mix
ERNMX-109	Crownvetch Seeding Mix (Naturalized)
ERNMX-110	Ernst Native Biomass Mix for Strip Mines & Natural Gas Production Sites
ERNMX-111	Ernst Native Habitat Mix for Strip Mines
ERNMX-181	Native Steep Slope Mix with Annual Ryegrass

THESE MIXES ARE GOOD FOR CONTROLLING EROSION AND PROVIDING FOOD AND/OR COVER FOR WILDLIFE.

VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See "Disclaimer," p. 11. For "Expectations of Native Species," see p. 8.



This wetland was constructed on a former mine site.

**SECOND & SUBSEQUENT GROWING SEASONS**

- › Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any material standing from the previous year close to the ground (approximately 2") on sites that are not too rough or steep. This will allow the soil to warm more quickly, stimulating emergence and growth of native plants and reducing the likelihood of shrub invasion.
- › If bioengineering materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Round-up®, Rodeo®, Garlon®, Garlon® 3A, Stinger®, or Mile-stone®. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.


**SPECIAL CIRCUMSTANCES**

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". Trimming should cease by mid-September. However, vegetation allowed to grow without mowing provides more protection for wildlife and aids in erosion control. ☼



Seedlings from a steep slope mix poking through an erosion control blanket.

# UTILITY RIGHT-OF-WAY SITES



This electricity transmission R-O-W is well-established with beneficial native grasses, such as *Panicum clandestinum* (Deertongue), and forbs, including various *Solidagos* (Goldenrods) and *Eupatorium fistulosum* (Joe Pye Weed).

**MANY UTILITY COMPANIES AND OIL AND GAS OPERATORS** recognize a unique opportunity to go a step beyond basic reclamation standards by using native species indigenous to the area being restored. In doing so, they are affirming the industry's commitment to environmental stewardship.



A pipeline seeded with a mix of native grasses, forbs, and legumes.

In the years after disturbance, affected areas can become biodiverse ecosystems with improved ecological function, greater wildlife populations, less erosion, and improved water and soil quality.

For example, a multiple-mile stretch of pipeline in the Marcellus and Utica shale plays may pass through wetlands, over steep mountain slopes, across rivers, and through agricultural areas and state game lands. These areas should be reclaimed with vegetation best matching the intended use of the land, the biodiversity that existed before disturbance, and with practices that best address such issues as erosion control, habitat fragmentation, and other environmental concerns. We can design a biodiverse native seed mix to mitigate the environmental impact and aid in ensuring regulatory compliance. We routinely work with environmental departments, consulting engineers, and contractors seeding a project. 🌱

For Upland Meadow Sites Establishment Guide, see p. 19; Disturbed Sites & Steep Slopes Establishment Guide, see p. 32; Wetland Sites Establishment Guide, see p. 38; Riparian Sites Establishment Guide, see p. 29.

Due to the diverse range of site types that most utility transmission lines and oil and gas pipelines traverse, we recommend reviewing the seed mixes found on our website. Our sales representatives will also be able to assist in identifying the most appropriate seed mix for the project and site type.



This former well pad is bringing ecological value back to the landscape.

# WET MEADOW & WETLAND SITES



*Pontederia cordata* (Pickerelweed) is a key native wetland plant that provides food for aquatic wildlife.

**WET MEADOWS & WETLAND SITES HAVE** soils made up of clay and high organic matter with high water tables or impervious layers that prevent drainage. These sites are wet most of the time. Wet Meadow examples: Roadside ditches, retention basins that catch run-off water (see p. 45), pond areas, or wetland edges. Wetland examples: Newly created wetlands and wetland restoration sites, retention basins with wetland functions, floodplains, pond edges, open water, or wet bioremediation sites.



Wetlands are massive nutrient filters that help clean our nation's water supply and protect rivers and oceans from pollution.



**HABITAT:**

Varies from partial shade to full sun; requires wet or saturated soil, standing water, or a high water table; generally populated with wetland and wet meadow species.



**FERTILITY:**

Due to the potential for water contamination, lime and fertilizer are not recommended; however, when topsoil has been depleted or removed, we recommend the addition of organic matter (compost). Check soil pH and select species adapted to that pH.



**SEEDING METHOD:**

Hand seed, broadcast seed, hydroseed, or drill seed when the water table is drawn down. It is not practical to seed any wetland where there is standing water or where severe flooding is likely to occur before germination. The same caution applies to mulching. Natural seed banks (seeds in wetland soils) often establish part of the vegetative cover.

**SITE PREPARATION**

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide, such as glyphosate (Rodeo®), triclopyr (Garlon® 3A), or a similar aquatic herbicide formulation, to control such undesirable vegetation as multiflora rose, honeysuckle, and woody species. CAUTION: Some persistent species, such as purple loosestrife, phragmites, Japanese knotweed, or reed canarygrass, may require multiple applications of glyphosate or triclopyr. Perennial weeds not addressed before establishment will be more difficult to remove later. These sites are often too wet to till. Newly constructed wetlands, retention basins, and wet construction sites should be seeded as soon after construction as possible. Leaving the surface rough by creating mounds and kettles for an undulating microtopography can be very beneficial in obligate wetlands.

**GROWING SEASON MAINTENANCE**

**FIRST GROWING SEASON**

- › When feasible (the ground isn't too slippery or mucky to safely walk), post-planting maintenance will provide the best results for wet meadows and wetlands. Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a string trimmer.



Trimming reduces competition by fast-growing weeds for sunlight and nutrients needed by slower growing perennial natives. Trimming should cease by mid-September.

- › Problem weeds should be hand pulled or spot sprayed with an approved aquatic herbicide, such as Rodeo® or Garlon® 3A.

**SECOND & SUBSEQUENT GROWING SEASONS**

- › Problem weeds, such as purple loosestrife, phragmites, Japanese knotweed, and reed canarygrass, should be hand pulled or spot sprayed with an approved aquatic herbicide, such as Rodeo® or Garlon® 3A. ✿

**WET MEADOWS & WETLAND SITES SEED MIXES**

<b>ERNMX-120</b>	OBL-FACW Perennial Food & Cover Wetland Mix
<b>ERNMX-122</b>	FACW Meadow Mix
<b>ERNMX-128</b>	Seasonally Flooded Wildlife Food Mix
<b>ERNMX-131</b>	OBL Wetland Mix
<b>ERNMX-137</b>	Specialized Wetland Mix for Shaded OBL-FACW Areas
<b>ERNMX-138</b>	Wildlife Food & Shelter Mix
<b>ERNMX-175</b>	Waterfowl Buffet Mix for Wetland Enhancement
<b>ERNMX-176</b>	Passive Acid Mine OBL Wetland Mix

**THESE MIXES ARE GOOD FOR VEGETATING DISTURBED OR MAN-MADE WETLANDS AND PROVIDE FOOD AND/OR COVER FOR INSECTS, BIRDS, AND MAMMALS.**

**VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.**

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See “Disclaimer,” p. 11. For “Expectations of Native Species,” see p. 8.

# PARTIALLY SHADED SITES



Woodland openings can be developed to provide crucial early successional habitat between old growth forests and adjoining grasslands.

**PARTIALLY SHADED SITES ARE** characterized as having a relative lack of direct sunlight, typically caused by trees and shrubs that reduce their exposure. Examples: Woodland openings and sites associated with bioengineering installations.



*Aster divaricatus* (White Wood Aster) is frequently found along wooded edges and provides great pollinator value in late summer and early fall.



**HABITAT:**

Typically in moderate shade; many native species are adapted to moderate shade and the protective habitat around trees; shade tolerant native grass species, such as *Agrostis perennans* (Autumn Bentgrass), *Chasmanthium laxum* (Slender Woodoats), *Cinna arundinacea* (Wood Reedgrass), *Elymus hystrix* (Bottlebrush Grass), *Elymus riparius* (Riverbank Wildrye), *Elymus virginicus* (Virginia Wildrye), and *Panicum clandestinum* (Deertongue), provide early protection for emerging herbaceous species. Note: For understory of longleaf pine plantings, high biomass producing species, such as switchgrass, big bluestem, and indiagrass, should be avoided. Fire can be too hot for longleaf pine seedlings or trees when these species are burned.



**FERTILITY:**

The addition of organic matter (compost) is most important. Check soil pH and select species adapted to that pH.



**SEEDING METHOD:**

Hand seed, broadcast seed, or hydroseed. Use a garden rake, drag, or roll the surface to incorporate the seed into the soil 1/4"-1/2" deep. A seed drill may be used when sufficient room exists for operation.


**SITE PREPARATION**

These sites generally involve working around trees and shrubs while minimizing damage to trunks and roots. Undesirable vegetation must be controlled by tilling or direct spraying with glyphosate. Invasive weeds not addressed before establishment will be difficult to remove later. The soil needs to be loosened in order to establish seed-to-soil contact and dense leaf litter should be broken up with a rototiller. Seedlings can emerge from light leaf litter if planted at the proper depth. Light mulch or hydromulch can protect the seeds and soil until germination. Seeding and mulching around bioengineering material should take place immediately after installation. If installation cannot take place immediately after grading, temporary seeding and mulching are recommended.

**GROWING SEASON MAINTENANCE**

**FIRST GROWING SEASON**

› Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a brush hog mower or string trimmer. Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients need-



*Apocynum cannabinum* (Indianhemp) is attractive to butterflies and songbirds.



*Carex scoparia* (Blunt Broom Sedge) provides food and cover for songbirds, ruffed grouse, and ducks.

ed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.

- › If bioengineering materials were used on the site, mowing should be above the new growth of these materials. Trimming should cease by mid-September.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Stinger®, or Milestone®. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after they have had two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

**SECOND & SUBSEQUENT GROWING SEASONS**

- › Prior to new spring growth reaching 2” (e.g., shortly after forsythia or redbud blooms), trim any material standing from the previous year close to the ground (approximately 2”). This will allow the soil to warm more quickly, stimulating emergence and growth of herbaceous native plants and reducing invasion of woody undergrowth. In certain ecosystems, controlled burning by certified professionals can achieve the same results.
- › If bioengineering materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.
- › Problem weeds should be hand pulled or spot sprayed. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after they have had two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

**SPECIAL CIRCUMSTANCES**

- › If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8”. Trimming should cease by mid-September. ✿

PARTIALLY SHADED SITES SEED MIXES	
ERNMX-132	Right-of-Way Non-Native Woods Mix
ERNMX-132-1	Right-of-Way Native Woods Mix with Annual Ryegrass
ERNMX-140	Partially Shaded Area Roadside Mix

THESE MIXES PROVIDE FOOD AND/OR COVER FOR WILDLIFE AND STABILIZATION OF SOILS IN PARTIALLY SHADED AREAS.

VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See “Disclaimer,” p. 11. For “Expectations of Native Species,” see p. 8.

# STORMWATER MANAGEMENT SITES



Mature rain garden at the Crawford County Fairgrounds in Meadville, Pennsylvania using ERNMX-180 Rain Garden Mix.

**STORMWATER MANAGEMENT FACILITY SITES ARE** generally a best management practice integrated throughout land development projects which provide for a volume of water storage, infiltration, and evaporation mimicking the natural rate of runoff or groundwater recharge.



The sites (e.g., size, location, and depth) must be designed and constructed according to all applicable ordinances and under the direction of a qualified design professional. In addition to planting trees and shrubs, seeding with native plant species is an economical way to vegetate stormwater management sites. Erosion control fabric, mulch, or hydromulch is necessary to control erosion during and after construction.

Examples: Detention basins - temporarily impound run-off water, allowing for release at controlled rates; retention basins - stormwater management facilities with permanent impoundment or pool for improving water quality; bioretention basins and rain gardens - dynamic living microbiological systems that enhance retention, infiltration, and evaporation of run-off water while remaining attractive to wildlife.

## PLANT SELECTION

In all stormwater management facilities, plants prevent erosion and slow water movement, hold or convert pollutants, enhance infiltration and evapotranspiration, and encourage wildlife. Plant species or mixes may be selected that meet critical objectives and extreme conditions of the site. Native grasses produce fibrous root systems that tolerate fast-moving water. Woody and herbaceous species add aesthetics, provide wildlife food and habitat, and assist with evapotranspiration while preventing erosion.

To avoid stand failure, select a seed mix appropriate for the site's hydrology (moisture status). If the site is chronically moist with long periods of inundation, a retention basin, FACW meadow, or OBL wetland mix is appropriate; if chronically moist with occasional inundation (immediately after a rainfall) and periodic drying out, a FACW meadow or riparian floodplain mix is appropriate; or, if typically dry except for a 12-72 hour period after a rainfall, a detention basin or rain garden mix is appropriate.

## SITE PREPARATION

Prior to planting the site, invasive species, particularly those adapted to wet conditions, should be removed or sprayed using an approved herbicide by a licensed spray technician. Perennial weeds not addressed before establishment will be difficult to remove later. Normal vegetation can be worked into the topsoil which should be stockpiled until the final grade has been established.

With specifications and dimensions, on-site construction of the berm and outlets must be executed carefully to main-



tain structural integrity. The infiltration and plant growth areas should be loose and friable (easy to crumble), high in organic matter, and completed without compaction by heavy equipment. An excavator may be used to dig and drop each area of the bottom soil in a loose manner. Lime or compost can then be incorporated. The excavation machine does not move over the finished surface thereby avoiding unnecessary compaction. Native vegetation can be planted or seeded over this uneven absorbent surface.

## SEEDING AND PLANTING METHODS

Seeding and planting should begin immediately upon completion of the structure while the soil is still friable and before weeds emerge. Plan seeding and planting before the basin is flooded or allow the basin to drain before seeding. Broadcast seed evenly over each unit by hand seeding or hydroseeding. Seeding rates are generally low (1/2 lb per 1,000 sq ft). If broadcasting seed, see Seeding Methods on p. 15. If the soil is dry, incorporate the seed into the soil with a garden rake. Oats or rye can provide temporary vegetation to protect the soil until permanent vegetation is established. Using such native species as *Elymus virginicus* (Virginia Wildrye) can create an intermediate cover that is succeeded by long-term native vegetation. Straw mulch or straw coconut mats are frequently used to control erosion and protect emerging seedlings from extreme temperatures and drying out. Mulch should be sparse to allow sunlight to reach the ground. If the site is a retention basin, refer to the Wet Meadow & Wetland Sites Establishment Guide, p. 38.

Transplanted seedlings and shrubs may need to be watered until they become well-rooted. Irrigating seeded areas is beneficial until seedlings become established.

## GROWING SEASON MAINTENANCE

### FIRST GROWING SEASON

- › Whenever canopy height (overall vegetation) reaches 18"-24", use a brush hog mower or string trimmer to trim the meadow to 8". Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.
- › If bioengineering or containerized woody materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Rodeo®, Garlon® 3A, or Milestone® (do not use Milestone® where standing water is present). Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after they have had two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.



An urban rain garden catch basin utilizing ERNMX-180 Rain Garden Mix.

**SECOND & SUBSEQUENT GROWING SEASONS**

- › Prior to new spring growth reaching 2” (e.g., shortly after forsythia or redbud blooms), trim any material standing from the previous year close to the ground (approximately 2”). If the seed mix included sedges, trimming should be no lower than 2” above the crowns that produced the prior year’s growth. This will allow the soil to warm more quickly, stimulating emergence and growth of native plants and reducing the likelihood of shrub invasion.
- › If bioengineering or containerized woody materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.
- › Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Rodeo®, Garlon® 3A, or Milestone®. Be vigilant in controlling vines or spiny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.

**SPECIAL CIRCUMSTANCES - SECOND GROWING SEASON**

- › If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8”. If bioengineering or containerized woody materials were used, trimming should be above or around new growth of the plants. Trimming should cease after mid-September.

**GENERAL MAINTENANCE**

- › In addition to structural maintenance, siltation should be removed as needed. Close mowing throughout the growing season or extensive chemical use is not conducive to water quality improvement and wildlife habitat. 🌱

STORMWATER MANAGEMENT SITES SEED MIXES	
ERNMX-126	Retention Basin Floor Mix - Low Maintenance
ERNMX-127	Retention Basin Wildlife Mix
ERNMX-128	Seasonally Flooded Wildlife Food Mix
ERNMX-154	Floodplain Mix
ERNMX-180	Rain Garden Mix
ERNMX-180-1	Rain Garden Grass Mix
ERNMX-180-2	Southeast Rain Garden Mix
ERNMX-183	Native Detention Area Mix

THESE MIXES ADDRESS STORMWATER. VISIT [ERNSTSEED.COM](http://ERNSTSEED.COM) FOR MORE OPTIONS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See “Disclaimer,” p. 11. For “Expectations of Native Species,” see p. 8.

# SOUTHEASTERN U.S. SITES



The attractive flowers and foliage of *Lespedeza virginica* (Slender Lespedeza) add color to native meadows and its bountiful seeds feed game birds.

**SOUTHEASTERN U.S. SITES INCLUDE** regions within Alabama, the Carolinas, southern Virginia, Georgia, Mississippi, and Florida. Sites in the Southeast typically have sandy or clay-rich soils subject to drought. These characteristics necessitate careful attention to timing and preparation. Examples: Coastal Plain soils, Piedmont, and sandy soils (refer to Upland & Meadow Sites Establishment Guide, p. 19).



A Florida meadow.



**HABITAT:**

Southeastern sites have a longer growing season; therefore, plants native or adapted to the region should be selected; planting from November to March is ideal (when possible) as temperatures are adequate and rain events are frequent; if irrigation is available, planting can continue into the later months of spring and early summer.



**FERTILITY:**

With the exception of organic matter, natural fertility is generally adequate. Check soil pH and, if necessary, add lime to achieve a pH of at least 6.0.



**SEEDING METHOD:**

Drill seeding is recommended; however, broadcast seeding is an alternative preceded by rolling or tracking.

## SITE PREPARATION

If the site was previously a lawn or crop field to which herbicides were applied, it is important to allow the appropriate time interval for the herbicide residues to break down prior to planting. Some herbicide residues can prevent seedlings from germinating.

Competition from invasive or undesirable vegetation is the most limiting factor in upland meadow preparation. Prior to planting, all such vegetation must be fully controlled. Typical control strategies include repeated tilling, smothering with black plastic, or herbicides. While any of these methods may control existing weeds, they will not kill all weed seeds lying dormant in the soil. Seeds of such species as velvetleaf and pokeweed may germinate many decades after the species last flowered on the site.

When using the tillage strategy, a site is disc harrowed every two weeks for one to two months. The underlying principle of this process is that the root system of perennial species will be worn out to the point of killing the species. In addition, tillage will stimulate germination of some dormant weed seeds that will be killed with subsequent tillage. Planting should not take place until perennial species are completely killed.

Black plastic may also be used to kill weeds. It may be laid across tilled or untilled soil and anchored down by burying the edges in soil or laying boards or bricks across the surface. This strategy should be utilized during a growing season when the intent is to fall plant the same year or spring plant the following year.



*Baptisia albescens* (Spiked Wild Indigo) in a South Carolina pine flatwoods.

Use of an approved herbicide, such as glyphosate (Roundup® or Rodeo®), by a licensed spray technician is the most common and least time-intensive method for the control of existing vegetation. Since herbicides are most effective on actively growing plant tissues, they are very effective on new growth in the spring. Spraying should begin when growth reaches 6". A follow-up spray application one to two weeks later will address skips or persistent species. If substantial plant tissue remains on the surface following a full kill by herbicides, a close mowing, tillage, or burning may be necessary to achieve good seed-to-soil contact.

To prevent reinfestation of some weed species, use of an appropriate selective herbicide in conjunction with a seed mix tolerant of that herbicide may be necessary.

#### **SANDY SOILS**

Sandy soils behave differently under cultivation than those containing clay. It is essential to plant seed 1/2" deep into a firm seedbed with a seed drill if possible (Eastern Gamagrass should be planted 1" deep). Truax and other similar drills can accommodate a variety of seeds and have been proven effective in the Southeast. High sand content in these soils makes broadcasting seed less effective due to poor seed-to-soil contact. Seedbeds should be firmed to where one does not sink past the sole of his/her shoe when walking the prepared site. Soil amendments may be added as necessary to maintain proper levels of organic matter and achieve a pH of at least 6.0.

#### **CLAY-RICH SOILS**

Without topsoil, soils containing high clay levels can be as hard as brick and pose a formidable challenge for successful cultivation. These soils are extremely low in organic matter which allows the small clay particles to settle and become compacted after a rain event. They are often iron-rich, leading to a distinctive red color. To prevent the clay from hardening after a rain from which seedlings cannot emerge, increase soil organic matter by incorporating 1"-2" of well-decomposed organic matter or compost and working it into the top-most soil prior to planting using a tiller, harrow, disc, or similar implement. Cultivating the top 6"-8" of soil will aid in root development of emergent seedlings and allow some percolation of rainwater that would otherwise run off the surface with little to no infiltration and carry the seed away with it. These initial preparations are critical for the successful establishment of native plants in this challenging soil. Since soil compaction is minimized, drilling seed 1/4"-1/2" deep is the preferred planting method. Even with additional organic matter, this clay-rich soil will compact easily; therefore, operating heavy equipment over the planted site should be avoided.



Quail-friendly planting featuring *Chamaecrista fasciculata* (Partridge Pea), *Panicum anceps* (Beaked Panicgrass), *Rudbeckia hirta* (Blackeyed Susan), and *Schizachyrium scoparium* (Little Bluestem).

## GROWING SEASON MAINTENANCE

Refer to Upland & Meadow Sites maintenance, p 19.

When spot spraying in soils with low organic matter and high sand levels, begin with lower than recommended concentrations of herbicides for weed control to avoid valuable crop burnout. Chemical breakdown of many herbicides is achieved via soil microbes that generally feed off organic material. With less organic material available in the soil, there will be a smaller population of microbes that may result in longer periods of exposure to the active ingredients in herbicides. \*

SOUTHEASTERN US SITES SEED MIXES	
ERNMX-169	Southeast Annual & Perennial Wildflower Mix
ERNMX-187	Southeastern U.S. Roadside Native Mix

THESE MIXES ARE USED IN WELL-DRAINED SOILS WITH FULL SUN AND PROVIDE FOOD AND/OR COVER FOR WILDLIFE. MEADOW AND WILDFLOWER MIXES PROVIDE FOOD FOR INSECTS, INCLUDING NATIVE POLLINATORS.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not.

See “Disclaimer,” p. 11. For “Expectations of Native Species,” see p. 8.

# BIOMASS PRODUCTION SITES



Calvin Ernst with a mature stand of *Panicum virgatum* (Switchgrass).

**BIOMASS PRODUCTION SITES** are sites planted with warm season grasses specifically for harvesting or grazing livestock on the biomass (vegetative matter). Special focus is given to the fact that these perennial crops may be harvested from the site for a decade or more without replanting.



Switchgrass grows through corn stubble, reducing its need for weed control.

## ADAPTING YOUR FIELD FOR BIOMASS PRODUCTION

Every field has unique characteristics to be considered when establishing perennial native biomass species. These include soil type, hydrology, pH, fertility, erosion/run-off potential, compaction, existing vegetative cover, previous crop history, and harvest methods.

While natural soil type cannot be changed, native warm season species can tolerate virtually any soil type. Switchgrass can survive in a wide range of soil moisture. As is the case with row crops or alfalfa, biomass productivity will be directly related to soil quality.

Switchgrass can tolerate soil pH of 5.0-8.0 but will produce well at 6.0. Soil pH below 6.0 should be corrected with the addition of lime according to soil test recommendations.

Soil fertility is a function of the available nutrients that can be used by the plant. Warm season grasses (switchgrass in particular) can be more productive at lower fertility levels than row crops or alfalfa. Soil tests are required to determine soil fertility levels. Fertility levels referred to as moderate are generally adequate for biomass production. Fertilizer is not recommended for soils with moderate fertility levels. Adding nitrogen in the second and subsequent years is recommended based on expected yields.

Fields having a history of good weed control are the easiest to convert to native warm season grasses, such as those planted in corn or soybeans. Fields in conventional hay or pasture are somewhat more difficult to seed and require Roundup® to kill the cool season grasses as well as minimum tillage to work thatch into the subsoil.

No till or minimum till are the most effective means of seeding new warm season grasses. A limiting factor of no till is surface crop residue that prevents proper seed-to-soil contact, shades the germinating seedlings, and/or creates a





nitrogen deficiency during decomposition. Each situation requires customized tillage and herbicide considerations.

Tilling a field going to warm season grasses corrects surface roughness and incorporates crop residue before planting. Field surface must be smooth enough to spray, plant, mow, and bale.

Fields with perennial or invasive weeds that have not been mowed during previous growing seasons are the most difficult to prepare for native grass establishment.

Perennial vegetation of grasses and broadleaf weeds must be controlled prior to planting. Mowing or burning existing vegetation will produce new vegetative growth. Roundup® and a systemic broadleaf herbicide can then be sprayed to effectively kill undesirable species. Identify weeds present and use label rates to control weeds with one or more applications. Once controlled, seeding like a conventional hayfield may proceed. ❁



# OBSTACLES TO A SUCCESSFUL MEADOW ESTABLISHMENT



Stand failure in central Pennsylvania caused by excess application of *Lolium multiflorum* (Annual Ryegrass).



## OBSTACLES TO A SUCCESSFUL MEADOW ESTABLISHMENT

Over the past 20 years, we have had the opportunity to visit sites or see photographic evidence of many native meadows that failed to establish successfully. As a result, we have found some repetitive themes relating to these failures - site conditions, weeds, excessive cover crop seeding rates, or use of an inappropriate cover or companion crop.

### POOR SITE CONDITIONS

- › Poor pre-plant weed control: Native species require bare ground to germinate and grow and will not establish well in a site already vegetated with weeds or lawn.
- › Presence of excessive organic matter used as mulch: Mulch prevents good seed-to-soil contact. A seed may germinate but its radicle (first root) may be unable to find moisture and the seedling dies. Microorganisms that decompose mulch or compost consume nitrogen, a nutrient essential to plant growth. This results in a nitrogen-deficient environment in which a seedling will grow poorly or not at all. If using compost, be certain it has decomposed to where the parent material is unrecognizable.
- › Soil compaction: If the tip of a crowbar or piece of rebar cannot be pressed 3" deep into the soil, there is compaction which will result in a very poor or non-existent stand. For highly compacted soils, consider broadcasting 1"-2" of well-decomposed compost across the site followed by rototilling the compost in 6" deep.
- › Herbicide residues, including pre-emergent herbicides, from the previous year: Generally occurs on sites that were previously lawns with a lawn service contract for weed control within 12 months of planting. Also problematic is when the site was a crop field to which herbicides, such as Atrazine (atrazine), Princep® (simazine), Milestone® (aminopyralid), Stinger® (clopyralid), Resolve® Q (rimsulfuron), Cimarron® (metsulfuron methyl), or Synchrony® XP (chlorimuron ethyl), were applied within 12 months of planting.
- › Excessively high or low soil pH: For many plant species native to eastern North America, availability of many nutrients essential to plant growth is limited in soils with pH less than 4.5 or greater than 7.5. Ideal pH is 5.5-7.0. When pH is outside this range, species tolerant of the site's pH should be chosen (pH can be raised with lime or lowered with sulfur).



Canada thistle in a wetland restoration site.

- › Drought within two to six weeks after planting: Seeds cannot germinate without water and seedlings do not grow/ survive without water. In drought-prone areas, we recommend planting between late October or when soil temperatures are less than 55°F (13°C) at 3" deep and in spring when forsythia or redbud blooms.
- › Erosion (on steep slopes): Failure to use erosion control blankets or toe them in at the top of the slope. Erosion is also a problem on slopes where final tracking of the soil was not perpendicular to the slope, resulting in the seed being washed down to the bottom of the slope.
- › Presence of crownvetch, sericea lespedeza, trefoil, tall fescue, bahiagrass, bermudagrass, white Dutch clover, alsike clover, bindweed, mile-a-minute, Japanese hops, kudzu, or invasive shrub species not controlled prior to planting or volunteered from dormant seed when the soil was prepared for planting. These species can smother desirable but slower growing perennial natives.
- › Use of borrowed topsoil infested with seeds of invasive species.
- › Wildlife: Geese can eat seeds, seedlings, and mature plants while deer can be devastating to some wildflowers. If up to five deer are regularly observed in the area, it may be wise to plant a deer-resistant mix. If the deer population is sufficient to make growing a garden or fruit trees difficult, it may be impossible to grow wildflowers on the site. The same is true if a deer population of 20 or more is regularly observed in the area.
- › Lack of proper maintenance: Letting annual ryegrass cover crop or weeds, such as foxtail or ragweed, smother native seedlings during the first full growing season. Avoid this by trimming the meadow to 8" whenever growth exceeds 18"-24".
- › Failure to control invasive or problem species prior to planting or after germination: If there are vines or spiny plants in the landscape that were not planted, be vigilant in controlling them.

#### **APPLICATION OF A COVER CROP AT AN EXCESSIVE RATE**

Issues with an excessive rate of cover crop have generally been confined to the use of annual ryegrass. Excessive annual ryegrass applied to sites planted with native species smothers growth of the native meadow. We have not had complaints when annual ryegrass was used at 10-12 lb per acre.




Green foxtail smothered native seedlings during the first full growing season of this planting.

#### USE OF AN INAPPROPRIATE COVER OR COMPANION CROP

We do not recommend the use of bahiagrass, bermudagrass, or tall fescue as cover or companion crops in our native meadow mixes. Bahiagrass or bermudagrass as a cover crop will be impossible to control prior to new growth of perennial native species and will likely smother out the native species when used as a companion crop.

We do not recommend the addition of the following legumes to native meadows as they also tend to take over and smother out native species: *Lespedeza cuneata* (Sericea Lespedeza), *Coronilla varia* (Crownvetch), *Lotus corniculatus* (Bird's Foot Trefoil), *Trifolium pratense* (Red Clover), or *Trifolium repens* (Ladino Clover). If already present on the site, control these species by spot spraying with Roundup® (glyphosate).

If a legume is desired in a meadow mix, we recommend such native legumes as *Baptisia alba* (White Wild Indigo), *Baptisia albescens* (Spiked Wild Indigo), *Baptisia australis* (Blue False Indigo), *Baptisia tinctoria* (Yellow False Indigo), *Chamaecrista fasciculata* (Partridge Pea), *Chamaecrista nictitans* (Sensitive Pea), *Desmodium canadense* (Showy Ticktrefoil), *Desmodium paniculatum* (Panicked Ticktrefoil), *Lespedeza capitata* (Round-head Lespedeza), *Lespedeza frutescens* (Shrubby Bushclover), *Lespedeza virginica* (Slender Bushclover), *Senna hebecarpa* (Wild Senna), and *Senna marilandica* (Maryland Senna).



An abundance of Canada geese can decimate seeds and seedlings.



#### WHAT IS THE APPROPRIATE COVER CROP FOR A MEADOW?

We recommend the following cover crops and seeding rates:

- › Grain Oats: 30 lb per acre; planted January through July (areas north of the Mason-Dixon Line), January through April (areas south of the Mason-Dixon Line).
- › Grain Rye: 30 lb per acre; planted August through December (areas north of the Mason-Dixon Line), September through December (areas south of the Mason-Dixon Line), and September through April (moist sites).
- › Annual Rye: 10-12 lb per acre; planted year-round (dry sites).
- › Brown Top Millet: 10 lb per acre; planted May through August (dry sites south of the Mason-Dixon Line).
- › Japanese Millet: 10 lb per acre; planted May through August (wet sites).

These seeding rates are based on our experience with native meadows as well as our desire to establish strong, individual native plants. Planting cover crops that are too aggressive or thick diminishes the long-term viability of the perennial meadow plants. We have concluded that annual small grains, such as oats and rye, are the best cover crops or companion crops to plant with native seedlings when there is a need. Grain cover crops can reduce competition from aggressive weeds because they grow quickly and reduce the potential for erosion by providing quick cover. We generally do not recommend annual ryegrass as it is too aggressive and volunteer seedlings can be persistent. When using annual ryegrass with native species, do not exceed 12 lb per acre.

#### PROBLEM WEEDS FOR UPLAND MEADOWS

In much of our market area, crabgrass, giant foxtail, green foxtail, and ragweed can smother a meadow in the establishment year. If overtaken by these weeds, use a brush hog mower or string trimmer to trim the meadow to 8". Trimming below 4" will kill seedlings of many native species. A lawn mower is not recommended. \*

# BIOMASS



Calvin Ernst demonstrates the height potential of *Panicum virgatum* (Switchgrass).



Ground switchgrass is gaining popularity as an effective, readily accessible, and inexpensive livestock and poultry bedding material.

## BIOMASS PRODUCTION

Ernst Conservation Seeds is one of the largest switchgrass seed producers in the country, having more than 30 years' experience in the establishment, management, and harvest of native warm season grass seed and biomass.

Switchgrass, as well as other native warm season grasses, has attracted much attention as a potential source of alternative energy and sustainable fiber due to the following:

- › Native warm season grasses are perennial.
- › Native warm season grasses thrive in marginal soil conditions too wet or dry for traditional crops.
- › Native warm season grasses require minimal nutrient input.
- › Native warm season grasses are efficient in converting sunlight to useable biomass.
- › Native warm season grasses have proven soil, water, air, and wildlife benefits.

Switchgrass biomass production can vary greatly from one region to another. It is important to select switchgrass varieties well-suited to the growing conditions of your area. Please contact us and we will be happy to make recommendations.

A mix of switchgrass varieties adapted to your area can better acclimate to seasonal variation and soil conditions than a single variety. Diverse genetic material will allow the overall stand to thrive in a wider range of moistures, soil types, disease pressures, and weather.

Commonly, mixes that include other native grasses, such as *Andropogon gerardii* (Big Bluestem), *Sorghastrum nutans* (Indiangrass), *Schizachyrium scoparium* (Little Bluestem), *Panicum amarum* (Coastal Panicgrass), and *Spartina spp.* (Cordgrass spp.), may create a biomass product that will satisfy many conservation program requirements while also being able to be marketed.



### **SELECTING THE RIGHT BIOMASS VARIETIES**

Our supply of switchgrass seed comes from various sources, including our licensing of the varieties produced by intensive breeding programs at numerous institutions and regional populations made available from USDA plant materials centers. The regional populations have minimal genetic improvement for general physical characteristics and have been adequate for decades for erosion control, wildlife plantings, and in the Conservation Reserve Program (CRP). The new varieties, including ‘Colony’, ‘Performer’, ‘BoMaster’, ‘Timber’, ‘Liberty’, ‘Independence’, ‘Shawnee’, and RC Chippewa, have significant yield improvements and were bred with a focus on forage and biomass production.

### **FORAGE**

As with several other native warm season grasses, switchgrass can produce high-quality forage. Used in a system of rotational grazing, switchgrass allows for robust growth during hot summer months. University of Tennessee findings suggest that the nutrient content of this forage can be as high as 16%-17% crude protein.

Ground switchgrass straw is experiencing increased use as a forage extender in livestock feeds in that it works to increase bulk and dilute protein in operations with sources of high-protein feed.

### **NUTRIENT RUN-OFF PREVENTION**

Switchgrass has extensive roots, growing as deep as 5'-6'. In addition to serving as a superior soil stabilizer in erosion control, switchgrass and its root system form a tremendous ecological filter, soaking up such nutrients as nitrogen and sequestering carbon dioxide. Use of switchgrass as a buffer or part of a riparian system between agricultural activity and watersheds is seen by many as one of the best methods for protecting these priceless resources.

### **POULTRY AND DAIRY BEDDING**

Numerous studies have shown that ground switchgrass is easy on the pads of chicken feet, highly absorbent, and may



represent a benefit over other beddings in the reduction of ammonia. From a cost perspective, producers can grow switchgrass on their own marginal ground, then harvest and process it for their own bedding uses. In addition to helping with noise, site, and water pollution control, switchgrass can aid in making areas of marginal ground productive by supplying sustainable bedding.

For information on preparing a field for the establishment and production of native biomass, please visit p. 52-54.

For more information, please consult the Native Biomass FAQs at [www.ernstseed.com](http://www.ernstseed.com). ❁



# BIOENGINEERING MATERIALS



Live stakes installed and sprouting new growth.



## SOIL BIOENGINEERING

Soil bioengineering is the term for using plant material to arrest and prevent slope and streambank failure and erosion. The roots and stems serve as structural and mechanical elements in a slope protection system. Live cuttings and rooted plants are embedded in the ground in various arrays to serve as soil reinforcements, hydraulic drains, and barriers to earth movement. Once established, this living material effectively controls several stabilization and erosion control problems by binding the soil with its root system and creating a natural vegetative cover. Bioengineered sites are self-repairing and have the advantage of blending with natural surroundings.

## BIOENGINEERING MATERIALS & STREAM RESTORATION TECHNIQUES

Ernst Conservation Seeds is an experienced producer of common and specialized live soil bioengineering materials. We understand the unique needs of bioengineering site construction. Our material is grown, processed, and delivered to minimize on-site installation labor and maximize survival and quick establishment.



**PLACING AN ORDER**  
Please call for a quote as each bioengineering job and material selection is unique to a project.



**HOW SOON CAN I GET MY BIOENGINEERING ORDER?**

Since we custom cut your order, we require at least two-weeks' notice when bioengineering orders are placed.

## HELPFUL TIPS FOR A BIOENGINEERING PROJECT

Ernst Conservation Seeds' bioengineering products are dormant live material. Therefore, if installation cannot take place immediately upon arrival at the site, these products must be stored properly. Place in a cool, wet place out of direct sunlight, such as under straw or burlap. Open any pallets, boxes, and plastic bags so the material can be watered thoroughly. Do not allow the material to dry out. Soaking before planting significantly increases survival and growth rate.


For best survivability, the material should be planted during the dormant season, November 1st-April 30th. We do not guarantee any of our bioengineering material from May 1st-October 31st.

Overseeding and mulching a completed bioengineered project with the appropriate seed mixes protects the soil surface from erosion while adding biodiversity to the site.

## THE FOLLOWING MIXES ARE EXCELLENT FOR THIS PURPOSE:

<b>ERNMX-137</b>	Specialized Wetland Mix for Shaded OBL-FACW Areas
<b>ERNMX-138</b>	Wildlife Food & Shelter Mix
<b>ERNMX-178</b>	Riparian Buffer Mix

# STREAMBANK DIAGRAM



## ERNST CONSERVATION SEEDS PRODUCES BIOENGINEERING MATERIALS IN THE FOLLOWING FAST-ROOTING SPECIES:

BOTANICAL NAME	COMMON NAME	PLANT TYPE
<i>Cephalanthus occidentalis</i>	Buttonbush	Native Shrub
<i>Cornus amomum</i>	Silky Dogwood	Native Shrub
<i>Cornus sericea</i>	Red Osier Dogwood	Native Shrub
<i>Salix amygdaloides</i>	Peachleaf Willow	Native Tree
<i>Salix discolor</i>	Pussy Willow	Native Tree
<i>Salix exigua ssp. interior</i>	Sandbar Willow	Native Shrub
<i>Salix lucida</i>	Shining Willow	Native Shrub
<i>Salix nigra</i>	Black Willow	Native Tree
<i>Salix purpurea</i>	Streamco Willow	Naturalized Shrub
<i>Salix sericea</i>	Silky Willow	Native Shrub
<i>Salix x cottetii</i>	'Bankers' Dwarf Willow	Naturalized Shrub
<i>Sambucus canadensis</i>	Elderberry	Native Shrub
<i>Viburnum dentatum</i>	Arrowwood	Native Shrub
<i>Viburnum lentago</i>	Nannyberry	Native Shrub

For more information on the species listed above, refer to Partially Shaded Sites, p. 41.

Live Stake and Branch Layering Cross-Sections courtesy of United States Department of Agriculture, Natural Resources Conservation Service (NRCS), Engineering Field Handbook, December 1996, Chapter 16, "Streambank and Shoreline Protection", pp. 16-13 and 16-20.



Special thanks to John McCullah, Salix Applied Earthcare, for allowing us to use the information in his Bio-Draw software. More information is available at [www.biodraw.com](http://www.biodraw.com).

## LIVE STAKES

Stakes are dormant, live woody cuttings of a species with the branches trimmed off. Live staking performs an important function in creating a root mat that stabilizes the soil by reinforcing and binding soil particles together. Stake establishment also improves aesthetics and provides a habitat for wildlife. Live stakes may be used on their own to secure other bioengineering materials or as an anchor for erosion control and geo-fabric. Stakes or poles may also be inserted or driven through openings in rock structures, such as gabions, riprap, and other retaining structures.

**INSTALLATION NOTES:** Install stakes during their dormancy (late fall to early spring). Do not allow the material to dry out. Soaking before planting significantly increases survival and growth rate. Drive a pilot hole into firm soil and plant at right angles (buds oriented up) with at least two-thirds of its length underground. Plant stakes randomly or 3'-6' apart on triangular spacing. Tamp the soil down around the cuttings before watering. Irrigation may be necessary if a long dry spell or hot weather is expected following installation.

**LIVE STAKE SIZES:** 1/4"-1" diameter; 2'-4' lengths.



## BRUSH (BRANCH) LAYERS

Brush (branch) layers are living branches placed on a terrace along the contours of a streambank and interspersed between layers of soil. This technique is used to repair a slump or gully and is most effective for revegetating scour holes.

**INSTALLATION NOTES:** Brush (branch) layers are placed on terraced benches with two-thirds of the basal material tilted into the slope and covered with soil. Branches should protrude beyond the face of the terrace. Before installing, soil terraces can be additionally protected by putting down geo-fabric. Starting at the bottom of the slope, secondary brush (branch) layers may be added every 3'-4' proceeding up the slope. Straw mulching the finished surface is recommended for moisture retention and additional erosion control. Planting should be during the dormant season.


**BRUSH LAYER SIZES:** 3 linear ft per bundle, 28-36 branches per bundle; available in 3'-6' bundles.

## BRUSH MATTRESSES

Brush mattresses are living branches layered 1-2 branches thick in a crisscross pattern on a streambank to form a living ground cover. The mattress formed protects the bank surface until the branches root and native vegetation is established. This living system normally roots in the entire bank face, encouraging natural infiltration and immediately acting as a sediment trap.

**INSTALLATION NOTES:** Place material with the basal ends located toward the bottom of the slope. Using 3' square spacing, drive dead wedge stakes into the plant material. Stretch wire or biodegradable twine or rope diagonally around the dead stakes and finish driving the stakes in to tighten the wire, twine, or rope and secure the mattress. Place a thin layer of soil over the entire area to encourage rooting. Irrigation is necessary immediately following installation.

**BRUSH MATTRESS SIZES:** 1/4"-3" diameter x random length x 1-2 or 3-4 branches thick.





### WATTLES/FASCINES

Wattles or fascines are living branches bound together in long tubular bundles. When placed in shallow trenches across the slope of a bank, these structures provide protection from erosion and create a sediment trap that provides immediate bank support even prior to root growth. Once established, this live rooting material grows into a living fence-like erosion barrier. Within one growing season, roots and shoots grow along the entire length of the structure and quickly stabilize the bank.

**INSTALLATION NOTES:** This technique is simple, effective, and installed with little site disturbance. Material is placed in 6" wide trenches on banks or slopes parallel to the stream contour and partially covered with soil. Wedge-like dead stakes secure them in place at 2'-3' intervals. Live stakes may also be used in conjunction with dead stakes to secure the material. Straw mulching the site following installation retains moisture and reduces surface erosion. Irrigation is necessary after installation if the soil is dry.

**WATTLE/FASCINE SIZES:** 6' or 8' lengths are recommended for ease of handling; available in the following diameters: 4"-5", 6"-8", 9"-12" (custom lengths and diameters are available).



### DEAD WEDGE STAKES

Dead wedge stakes are pieces of hardwood cut into long wedges to secure wattles, brush mattresses, and other applications of soil bioengineering and erosion control measures.

**STAKE SIZES:** 1-1/4" x 3-1/4" x 2-1/2' long.

### LIVE WHIPS

Whips are slender, live woody shrub material well-suited for very moist areas of stream edges, commonly used in conjunction with gabion structures, riprap, and geo-fabric.

**INSTALLATION NOTES:** Push whips into the ground as far as they will go without breaking. At least two-thirds of the whip should be covered with soil. Whips may be installed either by laying them on an angle or planting them erect in the soil. When using whips with hard structures, be sure they are long enough to reach into the soil and moisture behind or below the structure. (Example: If installing whips through riprap, consider the 3' depth; therefore, install a 6' whip at least 2' into the moist soil behind the stone and 1' above the surface of the riprap).

**LIVE WHIP SIZES:** 3/8"-1" diameter; 4'-6' lengths.



Herbaceous Perennial

***Acorus americanus***  
**SWEETFLAG**

**Native**  
Rhizomatous species; sometimes misnamed *Acorus calamus*, an introduced species; provides food and cover for wildlife.

**HABITAT:** Swamps, shallow pond water, wet meadows.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
pH: 5.6-7.2  
Bloom Period: Spring/Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 70,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Agrostis perennans***  
**AUTUMN BENTGRASS**

**Native**  
Rhizomatous species; adapted to dry areas, including steep slopes and forests with dappled to full sun; provides winter grazing for wildlife.

**HABITAT:** Dry open ground, areas in light shade.

**CHARACTERISTICS:**  
Height: Up to 3 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.5-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 8,000,000  
Seeding Rate: Up to 30 lb per acre alone; up to 2.4 lb per acre in a steep slope mix



Herbaceous Perennial

***Agrostis alba***  
**REDTOP**

**Naturalized**  
Rhizomatous sod-forming species; produces quick cover on road banks and diversion ditches for erosion control; adapts well to pipeline restoration; provides food for wildlife.

**HABITAT:** Coastal marshes, roadsides, open ground; establishes well in moist soils.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.5-8.0  
Bloom Period: Summer

Approx. Seeds Per Lb: 4,851,000  
Seeding Rate: Up to 25% of a mix; up to 40 lb per acre alone



Herbaceous Perennial

***Agrostis stolonifera***  
**CREEPING BENTGRASS**

**Naturalized**  
Stoloniferous sod-forming grass; used on reclamation sites, lawns, and golf course putting greens; also used for soil erosion control.

**HABITAT:** Wet meadows, shores, wet to dry fields, roadsides.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.1-7.5  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 6,130,000  
Seeding Rate: Up to 5% of a fine fescue mix; up to 40 lb per acre alone



Herbaceous Perennial


***Agrostis hyemalis***  
**WINTER BENTGRASS**

**Native**  
Bunchgrass; good for partially drained soils on moderately shaded roadsides; provides winter grazing for wildlife.

**HABITAT:** Roadsides, meadows, fields, moist to dry open, sterile soils.

**CHARACTERISTICS:**  
Height: Up to 2.5 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.0-7.5  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 8,500,000  
Seeding Rate: Up to 30 lb per acre alone; up to 2.4 lb per acre in a steep slope mix



Herbaceous Perennial

***Agrostis tenuis***  
**COLONIAL BENTGRASS**

**Naturalized**  
Rhizomatous species; provides erosion control; naturalized in many areas of the northeastern United States where sheep were grazed.

**HABITAT:** Cultivated in pastures, lawns, dry open ground, along roadsides.

**CHARACTERISTICS:**  
Height: Up to 2.2 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.9-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 6,130,000  
Seeding Rate: Up to 5% of a fescue mix; up to 40 lb per acre alone



***Alisma subcordatum***  
**MUD PLANTAIN**

**Native**  
Grows quickly in early spring; produces seed in the fall; ideal for vernal pools; provides food for pheasants and waterfowl.

**HABITAT:** Marshes, stream sides, muddy shores, pond margins, shallow water.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
Pollinator Value: Low  
pH: 5.0-7.0  
Bloom Period: Spring/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 825,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



***Andropogon gerardii***  
**BIG BLUESTEM**

**Native**  
Warm season bunchgrass; used for erosion control in sand and gravel pits, mine spoil, and on roadsides; contributes to diversified biomass production; high quality livestock forage; provides food and cover for wildlife.

**HABITAT:** Riverbanks, roadsides, meadows, open woods, savannas, tallgrass prairies.

**CHARACTERISTICS:**  
Height: Up to 6 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 6.0-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 144,000  
Seeding Rate: Up to 30% of a mix; up to 10 PLS lb per acre alone

**Herbaceous Perennial**



***Allium cernuum***  
**NODDING ONION**

**Native**  
Bulb-forming species; ideal ornamental for rock gardens; readily reseeds itself.

**HABITAT:** Dry rocky slopes, wood borders, rocky banks, prairies; often limestone outcrops.

**CHARACTERISTICS:**  
Height: Up to 2 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): White, Pink

Approx. Seeds Per Lb: 123,000  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Andropogon ternarius***  
**SPLITBEARD BLUESTEM**

**Native**  
Warm season bunchgrass; white seedheads stand out in upland habitats of the southern half of the United States.

**HABITAT:** Pinelands, sandhills, old fields.

**CHARACTERISTICS:**  
Height: Up to 3.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: High  
pH: 4.0-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 216,000  
Seeding Rate: Up to 10% of a mix; up to 10 PLS lb per acre alone

**Herbaceous Perennial**



***Alopecurus arundinaceus***  
**CREEPING FOXTAIL**

**Naturalized**  
Rhizomatous species; ideal for pipeline restoration where wildlife is desired; provides seed, forage, and cover for wildlife and domestic animals.

**HABITAT:** Wet hay meadows, margins of lakes and ponds, waterways.

**CHARACTERISTICS:**  
Height: Up to 3.9 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.5-8.4  
Bloom Period: Spring

Approx. Seeds Per Lb: 786,000  
Seeding Rate: Up to 10% of a mix

**Herbaceous Perennial**



***Andropogon virginicus***  
**BROOMSEDGE**

**Native**  
Warm season bunchgrass; historically used to make brooms; stiff straw stays erect on road cuts during the winter; provides cover for wildlife.

**HABITAT:** Pastures, open woods, old fields, hillsides, open ground with dry infertile soils.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.9-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 800,000  
Seeding Rate: Up to 2.5% of a mix; up to 10 PLS lb per acre alone

**Herbaceous Perennial**



**Herbaceous Perennial**

***Anthoxanthum odoratum***  
**SWEET VERNALGRASS**

**Naturalized**

Dense, clump-forming cool season bunchgrass; sweet fragrance of freshly mowed hay when crushed.

**HABITAT:** Open fields, meadows, roadsides; grows primarily in poor soils.

**CHARACTERISTICS:**

Height: Up to 2.4 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 738,000  
Seeding Rate: Up to 3% of a mix



**Herbaceous Perennial**

***Bromus inermis***  
**SMOOTH BROME**

**Naturalized**

Sod-forming cool season grass; provides food for livestock.

**HABITAT:** Roadsides; grows best in well-drained fine-textured soils.

**CHARACTERISTICS:**

Height: Up to 3.4 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.5-8.0  
Bloom Period: Summer

Approx. Seeds Per Lb: 143,000  
Seeding Rate: Up to 15% of a mix



**Herbaceous Perennial**

***Bouteloua curtipendula***  
**SIDEOATS GRAMA**

**Native**

Fast-emerging warm season bunchgrass for upland meadows where sight lines are important; used for surface mine revegetation, erosion control, and as a warm season companion crop; high winter forage value for wildlife and livestock.

**HABITAT:** Dry woods, dry calcareous clearings, dry prairies, sandhills.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.5-8.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 159,000  
Seeding Rate: Up to 35% of a mix; up to 10 PLS lb per acre alone



**Herbaceous Perennial**

***Calamagrostis canadensis***  
**CANADA BLUEJOINT**

**Native**

Rhizomatous grass; provides food and cover for deer, muskrats, and moose.

**HABITAT:** Swamps, wet meadows.

**CHARACTERISTICS:**

Height: Up to 4.6 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.5-8.0  
Bloom Period: Summer

Approx. Seeds Per Lb: 3,837,000  
Seeding Rate: Up to 0.5% of a mix; up to 2 PLS lb per acre alone



**Herbaceous Perennial**

***Bouteloua gracilis***  
**BLUE GRAMA**

**Native**

Fast-emerging warm season bunchgrass; used in dry highway medians, recreation area plantings, and in pure stands for erosion control; provides food for wildlife.

**HABITAT:** Dry prairies, sandhills.

**CHARACTERISTICS:**

Height: Up to 1.8 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 6.6-8.4  
Bloom Period: Summer

Approx. Seeds Per Lb: 724,000  
Seeding Rate: Up to 10% of a mix; up to 6 PLS lb per acre alone



**Herbaceous Perennial**

***Carex albolutescens***  
**GREENWHITE SEDGE**

**Native**

Tufted sedge; provides food for wildlife.

**HABITAT:** Low fields, meadows, marshes, floodplain forests, thickets.

**CHARACTERISTICS:**

Height: Up to 3.6 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.3-7.8  
Bloom Period: Spring/Summer

Seeding Rate: Up to 20% of a mix



**Carex comosa**  
**COSMOS SEDGE**

**Native**  
Bunch-type sedge; provides food and cover for wildlife.

**HABITAT:** Swamps, marshes, swales.

**CHARACTERISTICS:**  
Height: Up to 4 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.6-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 480,000  
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



**Carex granularis var. haleana**  
**LIMESTONE MEADOW SEDGE**

**Native**  
Bunch-type sedge; provides food for wildlife.

**HABITAT:** Wet meadows, swales.

**CHARACTERISTICS:**  
Height: Up to 2.7 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 6.0-7.2  
Bloom Period: Spring/Summer

Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



**Carex crinita**  
**FRINGED SEDGE**

**Native**  
Bunch-type sedge; provides food and cover for songbirds, ruffed grouse chicks, ducks, and moose.

**HABITAT:** Moist to wet woods, thickets, marshes, ditches, stream-banks.

**CHARACTERISTICS:**  
Height: Up to 4.8 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.0-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 720,000  
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



**Carex grayi**  
**GRAY'S SEDGE**

**Native**  
Bunch-type sedge; may be used for ornamental purposes in shaded areas; provides food and cover for wildlife.

**HABITAT:** Swamps, wet woods.

**CHARACTERISTICS:**  
Height: Up to 3 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 5.7-7.2  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 19,000  
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



**Carex frankii**  
**FRANK'S SEDGE**

**Native**  
Very rhizomatous sedge; establishes quickly from seed; provides food for wildlife.

**HABITAT:** Swamps, wet woods, streambanks, ditches.

**CHARACTERISTICS:**  
Height: Up to 2.6 ft.  
Minimum Root Depth: 9 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 5.9-7.2  
Bloom Period: Spring/Fall

Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



**Carex intumescens**  
**STAR SEDGE**

**Native**  
Bunch-type sedge; provides food and cover for wildlife.

**HABITAT:** Wet woods, meadows, swamps.

**CHARACTERISTICS:**  
Height: Up to 2.7 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Shade  
Drought Tolerance: No  
pH: 4.8-6.9  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 40,000  
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



**Carex lupulina**  
**HOP SEDGE**

**Native**  
Bunch-type sedge; ducks enjoy the large seed grains; provides food and cover for wildlife.  
**HABITAT:** Swamps, wet woods.  
**CHARACTERISTICS:**  
Height: Up to 3.9 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 6.2-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 94,700  
Seeding Rate: Up to 15% of a mix

**Herbaceous Perennial**



**Carex scoparia**  
**BLUNT BROOM SEDGE**

**Native**  
Bunch-type sedge; provides food and cover for songbirds, ruffed grouse chicks, ducks, and moose.  
**HABITAT:** Swamps, wet meadows, moist open ground.  
**CHARACTERISTICS:**  
Height: Up to 3 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Shade  
Drought Tolerance: No  
pH: 4.6-6.9  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,344,000  
Seeding Rate: Up to 15% of a mix

**Herbaceous Perennial**



**Carex lurida**  
**LURID SEDGE**

**Native**  
Bunch-type sedge; provides food and cover for songbirds, ruffed grouse chicks, ducks, and moose.  
**HABITAT:** Swamps, bogs, wet meadows, wet woods.  
**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.9-6.8  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 250,000  
Seeding Rate: Up to 20% of a mix

**Herbaceous Perennial**



**Carex shortiana**  
**SHORT'S SEDGE**

**Native**  
Bunch-type sedge; shortest sedge we grow; discovered by botanist Charles Wilkins Short; provides food and cover for wildlife.  
**HABITAT:** Calcareous wet meadows, rich woods, swamps.  
**CHARACTERISTICS:**  
Height: Up to 2.9 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: No  
pH: 4.7-6.9  
Bloom Period: Spring/Summer

Seeding Rate: Up to 10% of a mix

**Herbaceous Perennial**



**Carex pennsylvanica**  
**PENNSYLVANIA SEDGE**

**Native**  
Alternative stoloniferous sedge; tolerates sandy soils; ideal ground cover in a mature deciduous forest; may need prescribed fire to form a sod.  
**HABITAT:** Open woods, wooded slopes.  
**CHARACTERISTICS:**  
Height: Up to 1.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer

Seeding Rate: Available as plants only

**Herbaceous Perennial**



**Carex stipata**  
**AWL SEDGE**

**Native**  
Bunch-type sedge; matures early in the season; provides food and cover for wildlife.  
**HABITAT:** Wet meadows, swampy woods.  
**CHARACTERISTICS:**  
Height: Up to 3.7 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.9-7.9  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 544,000  
Seeding Rate: Up to 5% of a mix

**Herbaceous Perennial**



**Carex stricta**  
**TUSsock SEDGE**

**Native**  
Tussock-forming sedge; provides habitat for turtles; source of food and cover for wildlife.

**HABITAT:** Swamps, streambanks, wet meadows.

**CHARACTERISTICS:**

Height: Up to 4 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 3.5-7.0  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,800,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



**Chasmanthium laxum**  
**SLENDER WOODOATS**

**Native**  
Decorative bunchgrass; may be used on mesic to floodplain sites in shade to full sun.

**HABITAT:** Moist sandy soils of the Coastal Plain.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 4.5-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 85,000  
Seeding Rate: Up to 40% of a mix

**Herbaceous Perennial**



**Carex vulpinoidea**  
**FOX SEDGE**

**Native**  
Bunch-type sedge; often the earliest sedge to establish from seed; provides food and cover for wildlife.

**HABITAT:** Moist meadows, marshes, ditches.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 6.8-8.9  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,297,000  
Seeding Rate: Up to 35% of a mix

**Herbaceous Perennial**



**Cinna arundinacea**  
**WOOD REEDGRASS**

**Native**  
Bunch-type grass; excellent cover in forested wetlands; provides forage for wildlife.

**HABITAT:** Wet woods, swamps.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.0-8.5  
Bloom Period: Summer/Fall

Seeding Rate: Up to 10% of a mix

**Herbaceous Perennial**



**Chasmanthium latifolium**  
**RIVER OATS**

**Native**  
Decorative bunchgrass; adds variety and texture to wildflower mixes; great for riparian sites; provides food and cover for wildlife.

**HABITAT:** Riverbanks, alluvial woods.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 5.0-7.0  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 90,000  
Seeding Rate: Up to 40% of a mix

**Herbaceous Perennial**



**Cyperus esculentus**  
**CHUFA**

**Native**  
Rhizomatous species; tubers are eaten by wildlife that dig them from the soil.

**HABITAT:** Moist ground of fields, meadows, lawns, gardens.

**CHARACTERISTICS:**

Height: Up to 2.6 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.0-7.0  
Bloom Period: Summer/Fall

Seeding Rate: Up to 5 lb per acre in a mix

**Herbaceous Perennial**



**Herbaceous Perennial**

***Dactylis glomerata***  
**ORCHARDGRASS**

**Native**

Cool season bunchgrass with a dense root system; reliable grass for many grazing programs; provides excellent livestock forage.

**HABITAT:** Open fields, meadows, roadsides with well-drained medium-textured soils.

**CHARACTERISTICS:**

Height: Up to 4.1 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 5.0-7.5  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 427,000  
Seeding Rate: Up to 10 lb per acre alone



**Herbaceous Perennial**

***Elymus canadensis***  
**CANADA WILDRYE**

**Native**

Short-lived cool season bunchgrass; establishes quickly in disturbed areas; used for soil stabilization; provides food and cover for wildlife.

**HABITAT:** Alluvial thickets, stream-banks, meadows; establishes best in well-drained soils.

**CHARACTERISTICS:**

Height: Up to 3.8 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 5.0-7.9  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 114,000  
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



**Herbaceous Perennial**

***Deschampsia cespitosa***  
**TUFTED HAIRGRASS**

**Native**

Clump-forming grass.

**HABITAT:** Wet or boggy ground.

**CHARACTERISTICS:**

Height: Up to 3.9 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
pH: 3.5-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,100,000  
Seeding Rate: Up to 1.3% of a mix; up to 0.25 lb per acre in a wildflower mix



**Herbaceous Perennial**

***Elymus glabriflorus***  
**SOUTHEASTERN WILDRYE**

**Native**

Short-lived cool season bunchgrass; used for erosion control; provides good livestock forage; should not be mowed or grazed below 6".

**HABITAT:** Alluvial fields, clearings, floodplain forests.

**CHARACTERISTICS:**

Height: Up to 4.6 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Bloom Period: Summer

Seeding Rate: Up to 20% of a mix; up to 30 PLS lb per acre alone



**Herbaceous Annual**

***Echinochloa muricata***  
**BARNYARDGRASS**

**Native**

Excellent bunchgrass cover crop for moist and wet disturbed soils; *Echinochloa crusgalli* var. *frumentacea* (Japanese Millet) is a non-native substitute; provides food for songbirds and game birds.

**HABITAT:** Moist ground, alluvial shores.

**CHARACTERISTICS:**

Height: Up to 4.4 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall

Seeding Rate: Up to 10 lb per acre as a cover crop



**Herbaceous Perennial**

***Elymus hystrix***  
**BOTTLEBRUSH GRASS**

**Native**

Short-lived cool season bunchgrass; used for soil stabilization; provides food and cover for wildlife.

**HABITAT:** Dry to mesic forests, woods.

**CHARACTERISTICS:**

Height: Up to 4.3 ft.  
Shade Tolerance: Shade  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 75,000  
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

***Elymus riparius***  
**RIVERBANK WILDRYE**

**Native**  
Short-lived cool season bunchgrass; used for soil stabilization, often mixed with Virginia Wildrye; provides food and cover for wildlife.

**HABITAT:** Alluvial flats, meadows, streambanks, wet rich woods.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.5-7.2  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 125,000  
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

***Eragrostis hirsuta***  
**BIGTOP LOVEGRASS**

**Native**  
Bunchgrass; attractive addition to upland meadows; early fall color.

**HABITAT:** Open disturbed habitats, clearings, roadsides, fields, open woods.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Shade Tolerance: Full Sun  
Bloom Period: Summer/Fall

Seeding Rate: Up to 2% of a mix; up to 8 PLS lb per acre alone



Herbaceous Perennial

***Elymus virginicus***  
**VIRGINIA WILDRYE**

**Native**  
Short-lived cool season bunchgrass; used for soil stabilization and revegetation of wetlands, often found with Riverbank Wildrye; provides food and cover for wildlife.

**HABITAT:** Moist woods, meadows, riverbanks.

**CHARACTERISTICS:**  
Height: Up to 4.2 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 5.0-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 73,000  
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

***Eragrostis spectabilis***  
**PURPLE LOVEGRASS**

**Native**  
Low-growing, short-rhizomed bunchgrass; grows well in open areas; provides a short visual layer when used with little bluestem or sideoats grama; early fall color.

**HABITAT:** Sandy fields, pastures, roadsides, open woods, open areas; tolerates low fertility soils.

**CHARACTERISTICS:**  
Height: Up to 2.3 ft.  
Minimum Root Depth: 4 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.0-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 1,000,000  
Seeding Rate: Up to 2% of a mix; up to 8 PLS lb per acre alone



Herbaceous Perennial

***Eragrostis curvula***  
**WEeping LOVEGRASS**

**Naturalized**  
Fast-establishing warm season bunchgrass; grows south of the Mason-Dixon Line; used for soil stabilization on steep slopes.

**HABITAT:** Sandy roadsides, fields.

**CHARACTERISTICS:**  
Height: Up to 4.3 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.5-8.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,482,000  
Seeding Rate: Up to 5% of a mix; up to 3-5 PLS lb per acre alone



Herbaceous Perennial

***Festuca arundinacea***  
**TALL FESCUE**

**Naturalized**  
Cool season bunchgrass; used for mine reclamation, stabilizing grassed waterways, slopes, lawns, and recreation areas.

**HABITAT:** Low fertility acidic, clay, loamy, or sandy soils.

**CHARACTERISTICS:**  
Height: Up to 4.6 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
pH: 5.0-9.0  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 227,000  
Seeding Rate: Up to 35% of a mix; up to 300 lb per acre alone



Herbaceous Perennial

***Festuca ovina***  
**SHEEP FESCUE**

**Naturalized**

Fine-leaved cool season bunchgrass; adds texture to landscapes; used for reclamation, banks, and pastures.

**HABITAT:** Open woods, dry fields, roadsides.

**CHARACTERISTICS:**

Height: Up to 2 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Moderate  
Drought Tolerance: High  
pH: 5.5-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 565,000  
Seeding Rate: Up to 35% of a mix; up to 25 lb per acre in a wildflower mix; up to 220 lb per acre alone



Herbaceous Perennial

***Festuca rubra***  
**CREEPING RED FESCUE**

**Native**

Sod-forming species; used for a no-mow cover and erosion control; provides habitat for wildlife.

**HABITAT:** Dry woods, roadsides, open ground.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 5.0-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 454,000  
Seeding Rate: Up to 35% of a mix; up to 220 lb per acre alone



Herbaceous Perennial

***Festuca ovina var. duriuscula***  
**HARD FESCUE**

**Naturalized**

Fine-leaved cool season bunchgrass; used for reclamation and roadway revegetation where short vegetation is essential; good wildflower companion crop.

**HABITAT:** Low fertility well-drained soils.

**CHARACTERISTICS:**

Height: Up to 2 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.5-8.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 565,000  
Seeding Rate: Up to 35% of a mix; up to 25 lb per acre in a wildflower mix; up to 220 lb per acre alone



Herbaceous Perennial

***Festuca rubra ssp. commutata***  
**CHEWINGS FESCUE**

**Naturalized**


Fine-leaved fescue; used for erosion control on slopes, waterways, and reclamation areas.

**HABITAT:** Well-drained acidic soils.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 500,000  
Seeding Rate: Up to 35% of a mix; up to 220 lb per acre alone



Herbaceous Perennial

***Festuca ovina var. glauca***  
**BLUE FESCUE**

**Naturalized**

Fine-leaved, ornamental cool season bunchgrass; used for ground cover and erosion control.

**HABITAT:** Open woods, roadsides, dry fields.

**CHARACTERISTICS:**

Height: Up to 2 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer

Seeding Rate: Up to 35% of a mix; up to 25 lb per acre in a wildflower mix; up to 220 lb per acre alone



Herbaceous Perennial

***Glyceria grandis***  
**AMERICAN MANNAGRASS**

**Native**

Decorative, wetland cool season bunchgrass; provides food for waterfowl, muskrats, and deer.

**HABITAT:** Swamps, marshes, wet meadows, shallow water, brooksides.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 659,000  
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial

***Glyceria striata***  
**FOWL MANNAGRASS**

**Native**  
Rhizomatous cool season bunchgrass; stays green through winter; provides food for waterfowl, muskrats, and deer.

**HABITAT:** Marshes, wet woods, swamps, bogs.

**CHARACTERISTICS:**  
Height: Up to 3.7 ft.  
Minimum Root Depth: 4 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.0-8.0  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,540,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Juncus tenuis***  
**PATH RUSH**

**Native**  
Bunch-type grass; tolerates foot traffic but not close mowing; used for nest material.

**HABITAT:** Moist to dry, often heavily compacted, soils of woods, fields, paths.

**CHARACTERISTICS:**  
Height: Up to 2.6 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.5-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 29,000,000  
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Iris versicolor***  
**BLUEFLAG**

**Native**  
Rhizomatous species; provides food for waterfowl, marsh birds, and muskrats.

**HABITAT:** Wet meadows, bogs, marshes.

**CHARACTERISTICS:**  
Height: Up to 3.1 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Low  
pH: 5.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Blue, Purple

Approx. Seeds Per Lb: 18,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Koeleria macrantha***  
**JUNEGRASS**

**Native**  
Cool season bunchgrass; provides good spring forage for livestock and deer; provides food for small mammals and upland game birds.

**HABITAT:** Open woods, dry soils.

**CHARACTERISTICS:**  
Height: Up to 2 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Shade  
Drought Tolerance: High  
pH: 6.0-8.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 2,315,000  
Seeding Rate: Up to 1.5% of a mix



Herbaceous Perennial

***Juncus effusus***  
**SOFT RUSH**

**Native**  
Bunch-type grass with a wide geographic distribution; provides spawning grounds for bluegills in shallow water; source of food and cover for songbirds and waterfowl.

**HABITAT:** Swamps, moist fields, floodplains, shores, ditches.

**CHARACTERISTICS:**  
Height: Up to 3.8 ft.  
Minimum Root Depth: 24 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.5-8.8  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 45,359,000  
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Leersia oryzoides***  
**RICE CUTGRASS**

**Native**  
Very rhizomatous warm season grass; creates a natural sediment trap; not recommended for use in residential settings as the vegetation can cause cuts to the skin (which gives the species its common name); provides food for ducks and habitat for invertebrates.

**HABITAT:** Marshes, bogs, wet meadows.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.1-8.8  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 498,000  
Seeding Rate: Up to 5% of a mix



**Herbaceous Annual**

***Lolium multiflorum*  
ANNUAL RYEGRASS**

**Naturalized**

Short-lived cool season bunchgrass; provides quick protection against soil, wind, and water erosion; used as a companion or cover crop where erosion is an immediate concern; frequently reseeds itself in disturbed areas.

**HABITAT:** Loose fertile to semi-fertile soils.

**CHARACTERISTICS:**

Height: Up to 2.6 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.0-7.9  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 217,000  
Seeding Rate: Up to 10% of a turf mix; up to 12 lb per acre as a companion crop with natives



**Herbaceous Perennial**

***Panicum amarum*  
COASTAL PANICGRASS**

**Native**

Warm season bunchgrass; used for the stabilization of coastal dunes, wind erosion control, and reclamation of gravel and mine areas; provides food and cover for wildlife.

**HABITAT:** Sandy shores, dune grasslands.

**CHARACTERISTICS:**

Height: Up to 6.5 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 5.0-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 325,000  
Seeding Rate: Up to 30% of a mix; up to 8 PLS lb per acre alone



**Herbaceous Perennial**

***Lolium perenne*  
PERENNIAL RYEGRASS**

**Naturalized**

Fast-growing cool season bunchgrass; used for soil stabilization, reclamation, pastures, athletic fields, and lawns.

**HABITAT:** Medium fertility acidic, clay, or loamy soils.

**CHARACTERISTICS:**

Height: Up to 2.8 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.0-8.0  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 240,000  
Seeding Rate: Up to 70% of a mix; up to 300 lb per acre alone



**Herbaceous Perennial**

***Panicum anceps*  
BEAKED PANICGRASS**

**Native**

Bunchgrass; provides food and cover for wildlife; forage is of good value for cattle; seeds are eaten by upland birds.

**HABITAT:** Moist sandy soils of ditches, fields, savannas, low pine-lands.

**CHARACTERISTICS:**

Height: Up to 3.8 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall

Seeding Rate: Up to 30% of a mix; up to 10 PLS lb per acre alone



**Herbaceous Perennial**

***Onoclea sensibilis*  
SENSITIVE FERN**

**Native**

Rhizomatous fern; provides shelter for salamanders and frogs.

**HABITAT:** Marshes, swamps, moist open woods, wet meadows.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Fall

Seeding Rate: Up to 1% of a mix



**Herbaceous Perennial**

***Panicum clandestinum*  
DEERTONGUE**

**Native**

Warm season bunchgrass with a fibrous root system; excellent for erosion control and revegetation of acid mine spoil and pipelines through wooded areas; good component of a streambank stabilization mix.

**HABITAT:** Moist soils of woodland edges and clearings.

**CHARACTERISTICS:**

Height: Up to 4.2 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.0-7.5  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 350,000  
Seeding Rate: Up to 3% of a mix; up to 8 PLS lb per acre alone



***Panicum dichotomiflorum***  
**SMOOTH PANICGRASS**

**Native**  
Warm season bunchgrass; used as a cover crop for wetland establishment; readily reseeds itself; plants and seeds provide food for ducks, deer, rabbits, and muskrats.

**HABITAT:** Moist soils, open woods, meadows.

**CHARACTERISTICS:**

Height: Up to 3.7 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
pH: 4.8-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 392,000  
Seeding Rate: Up to 10% of a mix

**Herbaceous Annual**



***Peltandra virginica***  
**ARROW ARUM**

**Native**  
Bunch-type species; seeds must be stored cold and wet; seed pods containing numerous large seeds that ripen in the fall are a source of food for wood ducks, muskrats, and rails; foliage provides cover for aquatic mammals, wading birds, and waterfowl.

**HABITAT:** Swamps, stream or lake edges, tidal marshes.

**CHARACTERISTICS:**

Height: Up to 1.6 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: No  
pH: 5.0-8.8  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 600  
Seeding Rate: Up to 0.5 lb per 1,000 sq ft

**Herbaceous Perennial**



***Panicum rigidulum***  
**REDTOP PANICGRASS**

**Native**  
Bunchgrass; attractive seedheads and red foliage in late summer and early fall; common component of wetlands in the Southeast; palatable for livestock.

**HABITAT:** Wet soils of marshes, alluvial swamps, ditches, low woods.

**CHARACTERISTICS:**

Height: Up to 4.2 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.0-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 797,000  
Seeding Rate: Up to 35% of a mix

**Herbaceous Perennial**



***Poa palustris***  
**FOWL BLUEGRASS**

**Native**  
Cool season bunchgrass; establishes quickly in wetlands and retention basins; provides food and cover for wildlife.

**HABITAT:** Wet meadows, damp soils.

**CHARACTERISTICS:**

Height: Up to 4.5 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.9-7.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 1,900,000  
Seeding Rate: Up to 25% of a mix; up to 10 lb per acre in a wet meadow mix; up to 160 lb per acre alone (lawns)

**Herbaceous Perennial**



***Panicum virgatum***  
**SWITCHGRASS**

**Native**  
Rhizomatous warm season bunchgrass; used for biomass, soil stabilization on strip mine spoil and dikes, and in buffer strips for nutrient uptake; provides pasture and hay for cattle and sheep; source of food and cover for wildlife.

**HABITAT:** Brackish marshes, river-side prairies, open woods, prairies.

**CHARACTERISTICS:**

Height: Up to 6.1 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 4.5-8.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 259,000  
Seeding Rate: Up to 30% of a mix; up to 8 PLS lb per acre alone

**Herbaceous Perennial**



***Poa pratensis***  
**KENTUCKY BLUEGRASS**

**Naturalized**  
Rhizomatous, sod-forming cool season grass; used for lawns and pastures; provides food and cover for wildlife.

**HABITAT:** Well-drained lawns, open woods, roadsides, meadows; tolerates a wide range of soils.

**CHARACTERISTICS:**

Height: Up to 3.1 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.0-8.4  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,390,000  
Seeding Rate: Up to 30% of a mix; up to 160 lb per acre alone (lawns)

**Herbaceous Perennial**



**Herbaceous Perennial**

***Poa trivialis***  
**ROUGH BLUEGRASS**

**Naturalized**  
Cool season bunchgrass; good for use in detention basins; grows well in early spring; provides food and cover for wildlife.

**HABITAT:** Wet meadows, moist woods, roadsides.

**CHARACTERISTICS:**  
Height: Up to 3.4 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.8-7.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 2,500,000  
Seeding Rate: Up to 25% of a mix; up to 160 lb per acre alone (lawns)



**Herbaceous Perennial**

***Sagittaria latifolia***  
**DUCK POTATO**

**Native**  
Ornamental wetland species; seeds and large tubers provide food for waterfowl, songbirds, wading birds, muskrats, and beavers.

**HABITAT:** Swamps, wet shores, shallow water of ponds and streams.

**CHARACTERISTICS:**  
Height: Up to 4.4 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
Pollinator Value: Low  
pH: 4.7-8.9  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 67,000  
Seeding Rate: Up to 1% of a mix



**Herbaceous Perennial**


***Pontederia cordata***  
**PICKERELWEED**

**Native**  
Ornamental wetland bunchgrass; seeds must be stored cold and wet; provides food for wildlife.

**HABITAT:** Marshes, swampy edges of lakes and streams, along tidal shores.

**CHARACTERISTICS:**  
Height: Up to 3.4 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
Pollinator Value: High  
pH: 4.9-8.7  
Bloom Period: Spring/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 5,000  
Seeding Rate: Up to 10 lb per acre



**Herbaceous Perennial**

***Schizachyrium scoparium***  
**LITTLE BLUESTEM**

**Native**  
Long-lived bunchgrass; dense roots can grow to 8' deep; good for up-land meadows where sight lines are important; used for erosion control on droughty sites; provides summer forage for livestock; source of food and cover for wildlife; grows with open exposed surfaces that host ground nesting pollinators.

**HABITAT:** Old fields, roadsides, riverside prairies, open woods, slopes, meadows.

**CHARACTERISTICS:**  
Height: Up to 4.4 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 5.0-8.4  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 241,000  
Seeding Rate: Up to 30% of a mix; up to 10 PLS lb per acre alone



**Herbaceous Perennial**


***Puccinellia distans***  
**ALKALIGRASS**

**Naturalized**  
Bunchgrass; used for erosion control and along roadsides where salt runoff is prevalent.

**HABITAT:** Roadsides, disturbed sites.

**CHARACTERISTICS:**  
Height: Up to 2 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 1,200,000  
Seeding Rate: Up to 20% of a mix



**Herbaceous Perennial**

***Scirpus atrovirens***  
**GREEN BULRUSH**

**Native**  
Bunch-type species; provides cover for wildlife in wet areas.

**HABITAT:** Moist meadows, marshes, floodplain forests, ditches; tolerates clay and silt soils found in wet areas.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.0-8.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 11,300,000  
Seeding Rate: Up to 1% of a mix



***Scirpus cyperinus***  
**WOOLGRASS**

**Native**  
Bunch-type species; stands tall in marshes; provides food and cover for waterfowl and muskrats.  
**HABITAT:** Moist meadows, marshes, swamps, shores, ditches.  
**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.8-7.2  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 36,000,000  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Scirpus validus***  
**SOFTSTEM BULRUSH**

**Native**  
Rhizomatous species; provides spawning grounds for fish in shallow water; source of food for waterfowl and muskrats.  
**HABITAT:** Swamps, wet ditches, mud flats, pond and lake margins.  
**CHARACTERISTICS:**  
Height: Up to 8.1 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
pH: 5.4-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Red  
  
Approx. Seeds Per Lb: 496,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



***Scirpus expansus***  
**WOOD BULRUSH**

**Native**  
Very rhizomatous species; provides cover for wildlife.  
**HABITAT:** Marshes, wet meadows, swamps, swales.  
**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 10,000,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



***Sisyrinchium angustifolium***  
**NARROWLEAF BLUE EYED GRASS**

**Native**  
Early season food source for pollinators.  
**HABITAT:** Damp woods, floodplains, grassy places, meadows.  
**CHARACTERISTICS:**  
Height: Up to 1.6 ft.  
Minimum Root Depth: 4 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 5.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Blue  
  
Approx. Seeds Per Lb: 757,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



***Scirpus hattorianus***  
**NORTHERN BULRUSH**

**Native**  
Clump-forming species.  
**HABITAT:** Swamps, bogs, moist meadows, riverbanks, ditches.  
**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer

Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Sorghastrum nutans***  
**INDIANGRASS**

**Native**  
Attractive warm season bunchgrass; good for erosion control, landscaping, and roadside beautification; provides food and cover for wildlife.  
**HABITAT:** Riverside prairies, moist or dry fields, open woods, roadsides, serpentine barrens; grows best in deep well-drained soils.  
**CHARACTERISTICS:**  
Height: Up to 6 ft.  
Minimum Root Depth: 24 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 4.8-8.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 175,000  
Seeding Rate: Up to 30% of a mix; up to 8 PLS lb per acre alone

**Herbaceous Perennial**



**Herbaceous Perennial**

***Sparganium americanum***  
**EASTERN BUR REED**

**Native**  
Rhizomatous, emergent aquatic plant; provides food for waterfowl, muskrats, and beavers.

**HABITAT:** Muddy shores, shallow water rivers, streams, swamps, ponds.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: No  
pH: 4.9-7.3  
Bloom Period: Spring/Fall  
Flower Color(s): Green

Approx. Seeds Per Lb: 50,000  
Seeding Rate: Up to 20% of a mix



**Herbaceous Perennial**

***Sporobolus asper***  
**ROUGH DROPSEED**

**Native**  
Warm season bunchgrass; survives in very poor soil conditions with little organic matter.

**HABITAT:** Dry sandy soils.

**CHARACTERISTICS:**  
Height: Up to 3.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 5.5-7.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 760,000  
Seeding Rate: Up to 5% of a mix



**Herbaceous Perennial**

***Sparganium eurycarpum***  
**GIANT BUR REED**

**Native**  
Rhizomatous, emergent aquatic plant; provides food and cover for waterfowl, pheasants, muskrats, and beavers.

**HABITAT:** Bogs, swamps, lake margins, ditches, swampy meadows.

**CHARACTERISTICS:**  
Height: Up to 4 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: No  
pH: 5.0-8.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 1,500  
Seeding Rate: Up to 20% of a mix



**Herbaceous Perennial**

***Sporobolus cryptandrus***  
**SAND DROPSEED**

**Native**  
Early emerging warm season bunchgrass; provides food and cover for wildlife.

**HABITAT:** Dry sandy soils.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 6.6-8.0  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 5,600,000  
Seeding Rate: Up to 0.1% of a mix



**Herbaceous Perennial**

***Spartina pectinata***  
**PRAIRIE CORDGRASS**

**Native**  
Aggressive, rhizomatous warm season grass; root system provides erosion control on streambanks; source of cover for waterfowl, songbirds, and small mammals.

**HABITAT:** Marshes, wet prairies, shores, riverside prairies.

**CHARACTERISTICS:**  
Height: Up to 6.1 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 6.0-8.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 639,000  
Seeding Rate: Up to 5% of a mix; up to 8 PLS lb per acre alone



**Herbaceous Perennial**

***Sporobolus heterolepis***  
**PRAIRIE DROPSEED**

**Native**  
Decorative, fine-textured warm season bunchgrass; may be difficult to germinate; provides food and cover for wildlife.

**HABITAT:** Dry open ground.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
pH: 6.0-7.2  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 1,200,000  
Seeding Rate: Up to 3% of a mix




Herbaceous Perennial

**Tradescantia ohiensis**  
**OHIO SPIDERWORT**

**Native**  
Attractive clump-forming species; blooms open during the morning hours from spring through summer; flowers close in warm temperatures.

**HABITAT:** Meadows, prairies, thickets, dry rocky woodlands, floodplain forests.

**CHARACTERISTICS:**  
Height: Up to 4.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Spring/Summer  
Flower Color(s): Blue, Purple  
Approx. Seeds Per Lb: 128,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Tridens flavus**  
**PURPLETOP**

**Native**  
Warm season bunchgrass; adds late season color to native landscapes; used for soil stabilization; provides food and cover for wildlife.

**HABITAT:** Meadows, old fields, open woods, roadsides; tolerates low quality roadside and field soils.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.5-6.5  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 465,000  
Seeding Rate: Up to 15% of a mix; up to 10 PLS lb per acre alone drill seeded; up to 20 PLS lb per acre alone broadcast seeded



Herbaceous Perennial

**Tradescantia subaspera**  
**ZIGZAG SPIDERWORT**

**Native**  
Attractive clump-forming species; flowers close in warm temperatures.

**HABITAT:** Mesic to dry forests and woodlands.

**CHARACTERISTICS:**  
Height: Up to 4.1 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer  
Flower Color(s): Blue, Purple  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Tripsacum dactyloides**  
**EASTERN GAMAGRASS**

**Native**  
Warm season bunchgrass; plant in the fall as a dormant seeding with germination occurring in the spring; requires great patience as the seeds have high levels of dormancy; outstanding forage producer; provides food and cover for wildlife.

**HABITAT:** Riverside prairies, meadows, swamps, wet shores, open fields.

**CHARACTERISTICS:**  
Height: Up to 9.8 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.1-7.5  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 7,000  
Seeding Rate: Up to 50% of a mix; up to 20 PLS lb per acre alone



Herbaceous Perennial

**Tradescantia virginiana**  
**VIRGINIA SPIDERWORT**

**Native**  
Attractive, late spring clump-forming species; more compact than Ohio Spiderwort; flowers close in warm temperatures.

**HABITAT:** Prairies, mesic to dry upland forests, floodplain forests.

**CHARACTERISTICS:**  
Height: Up to 2.9 ft.  
Minimum Root Depth: 4 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 4.0-8.0  
Bloom Period: Spring/Summer  
Flower Color(s): Blue, Purple  
Approx. Seeds Per Lb: 175,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Annual

**Zizania aquatica**  
**WILDRIE**

**Native**  
Bunchgrass; seeds must be kept moist and require clean still water to grow; provides food for wood ducks, black ducks, and muskrats.

**HABITAT:** Tidal and non-tidal marshes that are 1" deep or more.

**CHARACTERISTICS:**  
Height: Up to 7.8 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
pH: 6.4-7.4  
Bloom Period: Summer/Fall

Approx. Seeds Per Lb: 11,000  
Seeding Rate: Up to 10 lb per acre when used with a mix; up to 40 lb per acre alone



Herbaceous Perennial

***Achillea millefolium***  
**COMMON YARROW**

**Naturalized**

Species with a long bloom period.  
**HABITAT:** Roadsides, fields, waste areas.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
pH: 6.0-8.0  
Bloom Period: Spring/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 2,852,000  
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

***Amsonia ciliata***  
**FRINGED BLUESTAR**

**Native**

Clump-forming species; provides food for native pollinators.

**HABITAT:** Sandhills, sandy woodlands.

**CHARACTERISTICS:**

Height: Up to 2.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring  
Flower Color(s): Blue

Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Agastache foeniculum***  
**ANISE (LAVENDER) HYSSOP**

**Native**

Provides food for pollinators.

**HABITAT:** Dry upland woods and prairies.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Blue

Approx. Seeds Per Lb: 1,540,000  
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

***Anemone canadensis***  
**CANADIAN ANEMONE**

**Native**

Very attractive rhizomatous species.

**HABITAT:** Sandy shores, thickets, damp prairies, wet meadows.

**CHARACTERISTICS:**

Height: Up to 2.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 131,660  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Agrimonia parviflora***  
**SMALL FLOWERED AGRIMONY**

**Native**

Adds structure and texture to wetlands.

**HABITAT:** Bogs, moist woods, thickets; tolerates poor soils.

**CHARACTERISTICS:**

Height: Up to 4.6 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 6.0-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 288,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Anemone virginiana***  
**THIMBLEWEED**

**Native**

Excellent for moderately shaded areas.

**HABITAT:** Dry open woods, slopes, thickets.

**CHARACTERISTICS:**

Height: Up to 3.8 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Low  
Bloom Period: Spring/Summer  
Flower Color(s): Green

Approx. Seeds Per Lb: 448,000  
Seeding Rate: Up to 1% of a mix



**Apocynum cannabinum**  
**INDIANHEMP**

**Native**  
Spreads by underground roots; used by Native Americans to make rope; may be allelopathic; stem fiber is used by songbirds and orioles to build nests; flowers are attractive to butterflies.

**HABITAT:** Woods, old fields, sandy flats, limestone bluffs, open ground.

**CHARACTERISTICS:**  
Height: Up to 3.8 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 4.5-7.0  
Bloom Period: Spring/Fall

Approx. Seeds Per Lb: 500,000  
Seeding Rate: Up to 0.3% of a mix

**Herbaceous Perennial**



**Asclepias syriaca**  
**COMMON MILKWEED**

**Native**  
Decorative species with a fragrance resembling a lilac; spreads from underground roots; essential food source for monarch butterfly caterpillars and other beneficial insects.

**HABITAT:** Fields, roadsides, open ground.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer  
Flower Color(s): Purple

Approx. Seeds Per Lb: 70,000  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



**Aquilegia canadensis**  
**EASTERN COLUMBINE**

**Native**  
One of the first flowers to bloom in the spring; nectar source for hummingbirds.

**HABITAT:** Cliffs, rocky slopes, dry woods; usually calcareous.

**CHARACTERISTICS:**  
Height: Up to 3.6 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Low  
pH: 5.0-8.0  
Bloom Period: Spring/Summer  
Flower Color(s): Red

Approx. Seeds Per Lb: 504,000  
Seeding Rate: Up to 1.3% of a mix

**Herbaceous Perennial**



**Asclepias tuberosa**  
**BUTTERFLY MILKWEED**

**Native**  
Showy clump-forming species with tuberous roots; essential food source for monarch butterfly caterpillars.

**HABITAT:** Dry woods, abandoned fields, roadsides, shale barrens; grows best in well-drained soils.

**CHARACTERISTICS:**  
Height: Up to 2.9 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
Pollinator Value: Very High  
pH: 4.8-6.8  
Bloom Period: Spring/Fall  
Flower Color(s): Orange

Approx. Seeds Per Lb: 70,000  
Seeding Rate: Up to 2.5% of a mix

**Herbaceous Perennial**



**Asclepias incarnata**  
**SWAMP MILKWEED**

**Native**  
Decorative rhizomatous species with a fragrance of bubble gum; essential food source for monarch butterfly caterpillars; we observe more chrysalis on this milkweed species than on any other in our production fields.

**HABITAT:** Swamps, floodplains, wet meadows.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
Pollinator Value: Very High  
pH: 5.0-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): Pink

Approx. Seeds Per Lb: 153,760  
Seeding Rate: Up to 3% of a mix

**Herbaceous Perennial**



**Aster divaricatus**  
**WHITE WOOD ASTER**

**Native**  
Rhizomatous species; widely distributed in wooded areas; provides food and habitat for wildlife.

**HABITAT:** Dry woods.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 670,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



Herbaceous Perennial

**Aster laevis**  
**SMOOTH BLUE ASTER**

**Native**  
Provides attractive late summer color in meadows; sought by deer for browse.  
**HABITAT:** Dry woods, rocky ledges, roadsides.  
**CHARACTERISTICS:**  
Height: Up to 6.3 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 5.8-7.8  
Bloom Period: Summer/Fall  
Flower Color(s): Blue, Purple  
Approx. Seeds Per Lb: 1,014,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Aster novae-angliae**  
**NEW ENGLAND ASTER**

**Native**  
Brilliant flowers stand out from a distance; provides food and cover for wildlife.  
**HABITAT:** Fields, roadsides, moist meadows.  
**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): Purple  
Approx. Seeds Per Lb: 1,100,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Aster lateriflorus**  
**CALICO ASTER**  
**Native**  
Attractive clump-forming species; provides food and cover for wildlife.  
**HABITAT:** Dry open places, open woods.  
**CHARACTERISTICS:**  
Height: Up to 4.1 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 5.2-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): White  
Approx. Seeds Per Lb: 800,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Aster oblongifolius**  
**AROMATIC ASTER**  
**Native**  
Longest blooming of our native asters, lasting up to and sometimes beyond two months; small oblong leaves have a lemony scent.  
**HABITAT:** Dry open places, calcareous hillsides, cliffs, bluffs.  
**CHARACTERISTICS:**  
Height: Up to 2.8 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): Blue, Purple  
Approx. Seeds Per Lb: 816,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Aster macrophyllus**  
**BIGLEAF ASTER**  
**Native**  
Showy rhizomatous species; source of color along wooded borders; provides habitat for wildlife.  
**HABITAT:** Woods, rocky slopes, woodland edges; grows best in fertile soils.  
**CHARACTERISTICS:**  
Height: Up to 4 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: Very High  
pH: 4.9-6.9  
Bloom Period: Summer/Fall  
Flower Color(s): Blue, Purple  
Approx. Seeds Per Lb: 800,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Aster pilosus**  
**HEATH ASTER**  
**Native**  
Aggressive rhizomatous species; establishes quickly on disturbed sites and rights-of-way.  
**HABITAT:** Dry fields, open woods, vacant lots, roadsides.  
**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 5.4-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): White  
Approx. Seeds Per Lb: 700,000  
Seeding Rate: Up to 2% of a mix



***Aster prenanthoides***  
**ZIGZAG ASTER**

**Native**  
Rhizomatous species; tolerates urban habitats.

**HABITAT:** Steambanks, wet meadows, low woods.

**CHARACTERISTICS:**  
Height: Up to 5.5 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 5.5-7.2  
Bloom Period: Summer/Fall  
Flower Color(s): Blue, Purple

Approx. Seeds Per Lb: 700,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



***Aster spectabilis***  
**SHOWY ASTER**

**Native**  
Attractive rhizomatous species; one of the prettiest asters to add to landscapes.

**HABITAT:** Woodland borders, pine barrens, clearings, roadsides, dry sandy soils.

**CHARACTERISTICS:**  
Height: Up to 2.9 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Purple, Blue

Approx. Seeds Per Lb: 630,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



***Aster puniceus***  
**PURPLESTEM ASTER**

**Native**  
Attractive rhizomatous species; adds color to FACW and OBL meadows; provides food for deer.

**HABITAT:** Swamps, wet meadows, riverbanks, moist roadsides.

**CHARACTERISTICS:**  
Height: Up to 7.1 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: Very High  
pH: 4.5-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Blue, Purple

Approx. Seeds Per Lb: 700,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



***Aster umbellatus***  
**FLAT TOPPED WHITE ASTER**

**Native**  
Decorative rhizomatous component of wet meadows.

**HABITAT:** Floodplains, swamps, moist woods, moist fields.

**CHARACTERISTICS:**  
Height: Up to 6.9 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 1,072,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



***Aster sagittifolius***  
**ARROWLEAF ASTER**

**Native**  
Rhizomatous species.  
**HABITAT:** Woodland edges, streambanks, open areas, roadside slopes.

**CHARACTERISTICS:**  
Height: Up to 4.4 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 1,487,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



***Baptisia alba***  
**WHITE WILD INDIGO**

**Native**  
Slow-developing legume; lasts for many years in low fertility soils.

**HABITAT:** Open upland woods, prairies.

**CHARACTERISTICS:**  
Height: Up to 4.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Spring/Summer  
Flower Color(s): White

Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial Legume**



**Herbaceous Perennial Legume**

**Baptisia australis**  
**BLUE FALSE INDIGO**  
**Native**  
 Historically used as a dye; may be used as a specimen plant.  
**HABITAT:** Open woods, riverbanks, sandy floodplains.  
**CHARACTERISTICS:**  
 Height: Up to 5 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: Medium  
 Bloom Period: Spring/Summer  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 22,000  
 Seeding Rate: Up to 0.5% of a mix



**Herbaceous Annual**

**Bidens aristosa**  
**SHOWY TICKSEED SUNFLOWER (BUR MARIGOLD)**  
**Native**  
 Reseeding annual in disturbed soils; provides aggressive first-year cover in wet meadows; seeds are eaten by wildlife.  
**HABITAT:** Meadows, fields, roadsides, ditches.  
**CHARACTERISTICS:**  
 Height: Up to 4.9 ft.  
 Minimum Root Depth: 8 in.  
 Shade Tolerance: Moderate  
 Drought Tolerance: Low  
 Pollinator Value: Medium  
 pH: 5.0-7.0  
 Bloom Period: Summer/Fall  
 Flower Color(s): Yellow  
 Approx. Seeds Per Lb: 130,000  
 Seeding Rate: Up to 2% of a mix



**Herbaceous Perennial Legume**

**Baptisia pendula**  
**LARGELEAF WILD INDIGO**  
**Native**  
 Rhizomatous early spring legume.  
**HABITAT:** Flatwoods, open woods, clearings.  
**CHARACTERISTICS:**  
 Height: Up to 4.9 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: Medium  
 Bloom Period: Spring/Summer  
 Flower Color(s): White  
 Seeding Rate: Up to 1% of a mix



**Herbaceous Annual**

**Bidens cernua**  
**NODDING BUR MARIGOLD**  
**Native**  
 Reseeding annual persistent in disturbed saturated soils; not for use in residential settings as seeds attach themselves to clothing; provides food and cover for wildlife.  
**HABITAT:** Swamps, wet shores, ditches.  
**CHARACTERISTICS:**  
 Height: Up to 3.3 ft.  
 Minimum Root Depth: 8 in.  
 Shade Tolerance: Moderate  
 Drought Tolerance: Low  
 Pollinator Value: Medium  
 pH: 5.1-7.0  
 Bloom Period: Summer/Fall  
 Flower Color(s): Yellow  
 Approx. Seeds Per Lb: 43,324  
 Seeding Rate: Up to 2% of a mix



**Herbaceous Perennial Legume**

**Baptisia tinctoria**  
**YELLOW FALSE INDIGO**  
**Native**  
 Essential food source for frosted elfin butterfly caterpillars.  
**HABITAT:** Dry sandy soils, clearings, dry open woods.  
**CHARACTERISTICS:**  
 Height: Up to 3.3 ft.  
 Minimum Root Depth: 16 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: High  
 Pollinator Value: Medium  
 pH: 5.8-7.0  
 Bloom Period: Spring/Fall  
 Flower Color(s): Yellow  
 Approx. Seeds Per Lb: 300,000  
 Seeding Rate: Up to 0.5% of a mix



**Herbaceous Annual**

**Bidens frondosa**  
**BEGGARTICK**  
**Native**  
 Reseeding annual in disturbed wetlands; not for use in residential settings as seeds attach themselves to clothing; provides food and cover for wildlife and ducks.  
**HABITAT:** Moist open ground, streambanks, roadsides.  
**CHARACTERISTICS:**  
 Height: Up to 4.1 ft.  
 Minimum Root Depth: 8 in.  
 Shade Tolerance: Moderate  
 Drought Tolerance: Low  
 Pollinator Value: Medium  
 pH: 5.2-7.2  
 Bloom Period: Summer/Fall  
 Flower Color(s): Yellow, Orange  
 Approx. Seeds Per Lb: 195,300  
 Seeding Rate: Up to 1.3% of a mix



***Centaurea cyanus*  
CORNFLOWER (BACHELOR'S  
BUTTON)**

**Naturalized**

Blooms in early spring if sown in the fall.

**HABITAT:** Meadows, flower beds.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Fall  
Flower Color(s): Blue

Approx. Seeds Per Lb: 90,000  
Seeding Rate: Up to 11% of a mix

**Herbaceous Annual**



***Cheiranthus allionii*  
WALLFLOWER**

**Naturalized**

Decorative early blooming component in wildflower mixes.

**HABITAT:** Fields, meadows.

**CHARACTERISTICS:**

Height: Up to 1.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring  
Flower Color(s): Yellow, Orange, Red

Approx. Seeds Per Lb: 345,000  
Seeding Rate: Up to 3% of a mix

**Herbaceous Biennial/  
Perennial**



***Chamaecrista fasciculata*  
PARTRIDGE PEA**

**Native**

Bunch-type, readily reseeding annual in disturbed upland sites; foliage is nutritious but can be poisonous and should be considered potentially dangerous to cattle; fruit and seeds can irritate the digestive tract of livestock; seeds are a source of food for quail.

**HABITAT:** Riverbanks, sandy soils, clearings, roadsides.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 5.5-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 65,000  
Seeding Rate: Up to 3% of a mix

**Herbaceous Annual Legume**



***Chelone glabra*  
TURTLEHEAD**

**Native**

Showy wetland species browsed by wildlife; produces nectar containing catapol which aids in reducing parasite loads in bumblebees.

**HABITAT:** Streambanks, wet woods, wet meadows, swamps.

**CHARACTERISTICS:**

Height: Up to 3.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 1,472,000  
Seeding Rate: Up to 0.3% of a mix

**Herbaceous Perennial**



***Chamaecrista nictitans*  
SENSITIVE PEA**

**Native**

Tap-rooted legume; can help to increase nitrogen availability in some soils; fruit and seeds can irritate the digestive tract of livestock; seeds provide food for quail, doves, and turkey; attracts parasitoid wasps that control plant-eating insects.

**HABITAT:** Fields, roadsides, clearings, near rivers, woodlands.

**CHARACTERISTICS:**

Height: Up to 1.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 206,570  
Seeding Rate: Up to 1% of a mix

**Herbaceous Annual Legume**



***Chrysanthemum leucanthemum*  
OXEYE DAISY**

**Naturalized**

Decorative bunch-type species; persistent on low fertility sites.


**HABITAT:** Fields, woods, meadows, roadsides.

**CHARACTERISTICS:**

Height: Up to 2.5 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
pH: 5.2-7.0  
Bloom Period: Spring/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 200,000  
Seeding Rate: Up to 2.5% of a mix

**Herbaceous Perennial**



2 mm Grid

Herbaceous Perennial

***Chrysanthemum maximum***  
**SHASTA DAISY**

**Naturalized**  
Attractive component of naturalized meadow mixes.

**HABITAT:** Gardens, meadows.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 400,000  
Seeding Rate: Up to 2.5% of a mix



Herbaceous Annual

***Coreopsis basalis***  
**GOLDENMANE TICKSEED**

**Native**  
Adds early and mid-season color to southeastern landscapes.

**HABITAT:** Fields and roadsides in sandy soils.

**CHARACTERISTICS:**

Height: Up to 3.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Spring/Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 500,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Cichorium intybus***  
**BLUE CHICORY**

**Naturalized**  
Attractive bunch-type species; persistent on roadsides and in compacted soils; blooms last all day.

**HABITAT:** Fields, roadsides, waste ground.

**CHARACTERISTICS:**

Height: Up to 3.4 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 6.0-7.5  
Bloom Period: Spring/Fall  
Flower Color(s): Blue

Approx. Seeds Per Lb: 426,400  
Seeding Rate: Up to 2.5% of a mix



Herbaceous Perennial

***Coreopsis grandiflora***  
**LARGEFLOWER TICKSEED**

**Native**  
Showy component in wildflower meadows.

**HABITAT:** Roadsides, upland woods.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
Pollinator Value: Medium  
Bloom Period: Spring/Summer  
Flower Color(s): Yellow

Seeding Rate: Up to 3% of a mix



Herbaceous Perennial Vine

***Clematis virginiana***  
**VIRGIN'S BOWER**

**Native**  
Hardy vine to include in wetland margins; attractive when in bloom and when seed is ripening.

**HABITAT:** Thickets, streambanks, low woods.

**CHARACTERISTICS:**

Height: Up to 16.3 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 5.0-6.8  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 192,000  
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

***Coreopsis lanceolata***  
**LANCELEAF COREOPSIS**

**Native**  
Popular for wildflower meadows and along roadsides.

**HABITAT:** Soils, thickets, fields, clearings, roadsides.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 6.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 221,000  
Seeding Rate: Up to 3% of a mix



**Coreopsis tinctoria**  
**PLAINS COREOPSIS**

**Native**  
Showy annual; produces flowers in a short period of time on low fertility sites.

**HABITAT:** Fields, meadows, roadsides, occasionally escaping to yards.

**CHARACTERISTICS:**  
Height: Up to 3.6 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 5.2-7.8  
Bloom Period: Spring/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 3,222,000  
Seeding Rate: Up to 1% of a mix

Herbaceous Annual



**Cosmos bipinnatus**  
**COSMOS**

**Naturalized**  
Robust with intense color; ideal for showy roadside plantings; attracts butterflies.

**HABITAT:** Disturbed sites, fields, roadsides.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Mixed

Approx. Seeds Per Lb: 72,000  
Seeding Rate: Up to 13% of a mix

Herbaceous Annual



**Coreopsis tripteris**  
**TALL COREOPSIS**

**Native**  
Long-lived species; very tolerant of competition.

**HABITAT:** Old fields, thickets, woodland edges, roadsides, moist low places.

**CHARACTERISTICS:**  
Height: Up to 8.1 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 200,000  
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



**Cosmos sulphureus**  
**SULPHUR COSMOS**

**Naturalized**  
Attractive summer color lasts into the fall.

**HABITAT:** Gardens, roadsides, other disturbed habitats.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 60,000  
Seeding Rate: Up to 15% of a mix

Herbaceous Annual



**Coronilla varia**  
**CROWN VETCH**

**Naturalized**  
Rhizomatous legume; good for controlling erosion on steep, dry rocky slopes; seeds are persistent in soils; can be aggressive in natural areas.

**HABITAT:** Rocky slopes, shallow, well-drained low fertility soils.

**CHARACTERISTICS:**  
Height: Up to 2.6 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: High  
pH: 5.0-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Pink

Approx. Seeds Per Lb: 140,000  
Seeding Rate: Up to 50% of a mix; up to 8 PLS lb per acre alone

Herbaceous Perennial Legume



**Dalea candida**  
**WHITE PRAIRIE CLOVER**

**Native**  
Supplies nitrogen to meadow soils; produces nutritious forage for deer; seeds are eaten by birds.

**HABITAT:** Dry prairies, dry upland woods.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: No  
Bloom Period: Spring/Summer  
Flower Color(s): White


Approx. Seeds Per Lb: 278,000  
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial Legume



***Dalea purpurea***  
**PURPLE PRAIRIE CLOVER**  
**Native**  
 Fixes nitrogen in prairie and meadow soils; excellent forage for wildlife.  
**HABITAT:** Dry prairies, open glades.  
**CHARACTERISTICS:**  
 Height: Up to 3.1 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 pH: 6.0-8.0  
 Bloom Period: Spring/Summer  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 300,000  
 Seeding Rate: Up to 1% of a mix

Herbaceous Perennial  
Legume



***Desmanthus illinoensis***  
**ILLINOIS BUNDLEFLOWER**  
**Native**  
 Legume with a deep taproot; fixes nitrogen in prairie and meadow soils that can be used by plants.  
**HABITAT:** Prairies, roadsides, meadows, riverbanks.  
**CHARACTERISTICS:**  
 Height: Up to 6 ft.  
 Minimum Root Depth: 12 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Moderate  
 pH: 5.0-8.0  
 Bloom Period: Summer  
 Flower Color(s): White  
 Approx. Seeds Per Lb: 85,000  
 Seeding Rate: Up to 1% of a mix

Herbaceous Perennial  
Legume




***Daucus carota***  
**QUEEN ANNE'S LACE**  
**Naturalized**  
 Persistent in meadows with poor soils; attracts butterflies.  
**HABITAT:** Roadsides, old fields, gardens, open ground.  
**CHARACTERISTICS:**  
 Height: Up to 4.6 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Bloom Period: Spring/Fall  
 Flower Color(s): White  
 Seeding Rate: Up to 1% of a mix

Herbaceous Biennial



***Desmodium canadense***  
**SHOWY TICKTREFOIL**  
**Native**  
 Nitrogen-fixing legume; not for use in residential settings as seeds attach themselves to clothing; seeds provide food for ground birds and small animals; attracts native pollinators.  
**HABITAT:** Open woods, meadows, thickets, riverbanks.  
**CHARACTERISTICS:**  
 Height: Up to 4.9 ft.  
 Shade Tolerance: Moderate  
 Drought Tolerance: Unknown  
 Bloom Period: Summer/Fall  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 72,500  
 Seeding Rate: Up to 2% of a mix

Herbaceous Perennial  
Legume



***Delphinium ajacis***  
**ROCKET LARKSPUR**  
**Naturalized**  
 Attractive tap-rooted species.  
**HABITAT:** Gardens, roadsides, other disturbed habitats.  
**CHARACTERISTICS:**  
 Height: Up to 2.9 ft.  
 Shade Tolerance: Moderate  
 Drought Tolerance: No  
 Bloom Period: Spring/Fall  
 Flower Color(s): Blue, White, Pink  
 Approx. Seeds Per Lb: 140,000  
 Seeding Rate: Up to 5% of a mix

Herbaceous Annual



***Desmodium paniculatum***  
**PANICLEDLEAF TICKTREFOIL**  
**Native**  
 Nitrogen-fixing legume; not for use in residential settings as seeds attach themselves to clothing; seeds provide food for wildlife; attracts native pollinators.  
**HABITAT:** Dry woods, fields.  
**CHARACTERISTICS:**  
 Height: Up to 3.7 ft.  
 Minimum Root Depth: 6 in.  
 Shade Tolerance: Moderate  
 Drought Tolerance: Moderate  
 Pollinator Value: Low  
 pH: 6.0-7.0  
 Bloom Period: Summer/Fall  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 200,000  
 Seeding Rate: Up to 1% of a mix

Herbaceous Perennial  
Legume



Herbaceous Perennial

***Dianthus barbatus***  
**SWEETWILLIAM**

**Naturalized**

Food source for hummingbirds and butterflies.

**HABITAT:** Gardens, roadsides, other disturbed habitats; grows best in fertile, moist well-drained soils.

**CHARACTERISTICS:**

Height: Up to 1.9 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer  
Flower Color(s): Red, Pink, White

Approx. Seeds Per Lb: 440,000  
Seeding Rate: Up to 2.5% of a mix



Herbaceous Perennial

***Eryngium yuccifolium***  
**RATTLESNAKE MASTER**

**Native**

Creates visual texture in native meadows; essential food source for the rare rattlesnake master borer moth.

**HABITAT:** Moist woods, moist or dry sandy soils, meadows, barrens.

**CHARACTERISTICS:**

Height: Up to 4.7 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 178,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Echinacea pallida***  
**PALE PURPLE CONEFLOWER**

**Native**

Attractive in wildflower meadows.

**HABITAT:** Dry open spaces.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 6.5-7.2  
Bloom Period: Summer  
Flower Color(s): Pink, Purple

Approx. Seeds Per Lb: 106,000  
Seeding Rate: Up to 7% of a mix



Herbaceous Perennial

***Eupatorium coelestinum***  
**MISTFLOWER**

**Native**

Rhizomatous, low-growing, late fall flower in wetland margins.

**HABITAT:** Floodplain forests, old fields, meadows, streambanks.

**CHARACTERISTICS:**

Height: Up to 3.1 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 5.5-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Blue

Approx. Seeds Per Lb: 1,500,000  
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

***Echinacea purpurea***  
**PURPLE CONEFLOWER**

**Native**

Source of herbal remedy; attractive species in meadows and along roadsides.

**HABITAT:** Open woods, open meadows, prairies, roadsides; grows best in moist well-drained soils.

**CHARACTERISTICS:**

Height: Up to 5.9 ft.  
Minimum Root Depth: 24 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: High  
pH: 6.5-7.2  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 115,664  
Seeding Rate: Up to 7% of a mix



Herbaceous Perennial

***Eupatorium fistulosum***  
**JOE PYE WEED**

**Native**

Showy, rhizomatous, hollow-stemmed Joe Pye Weed.

**HABITAT:** Floodplains, meadows, moist thickets, roadsides.

**CHARACTERISTICS:**

Height: Up to 7.7 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: High  
pH: 4.5-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Pink, Purple

Approx. Seeds Per Lb: 2,000,000  
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

***Eupatorium maculatum***  
**SPOTTED JOE PYE WEED**

**Native**  
Source of food for pollinators.  
**HABITAT:** Floodplains, swamps, alluvial thickets.  
**CHARACTERISTICS:**  
Height: Up to 6.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Purple  
Approx. Seeds Per Lb: 1,440,000  
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

***Euthamia graminifolia***  
**GRASSLEAF GOLDENROD**

**Native**  
Rhizomatous species; provides food and cover for wildlife.  
**HABITAT:** Wet meadows, riparian areas; tolerates poor soils.  
**CHARACTERISTICS:**  
Height: Up to 4.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow  
Approx. Seeds Per Lb: 5,600,000  
Seeding Rate: Up to 0.2% of a mix



Herbaceous Perennial

***Eupatorium perfoliatum***  
**BONESET**

**Native**  
Beautiful, hardy component of wetlands; seeds are eaten by swamp sparrows.  
**HABITAT:** Floodplains, swamps, bogs, streambanks, wet meadows.  
**CHARACTERISTICS:**  
Height: Up to 5.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: No  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): White  
Approx. Seeds Per Lb: 2,880,000  
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

***Gaillardia aristata***  
**PERENNIAL GAILLARDIA (BLANKETFLOWER)**

**Native**  
Attractive bunch-type, daisy-like flower for meadows and along roadsides.  
**HABITAT:** Plains, prairies, meadows, along roadsides.  
**CHARACTERISTICS:**  
Height: Up to 2.2 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.5-7.9  
Bloom Period: Summer  
Flower Color(s): Yellow  
Approx. Seeds Per Lb: 186,436  
Seeding Rate: Up to 10% of a mix



Herbaceous Perennial

***Eupatorium purpureum***  
**PURPLE NODE JOE PYE WEED**

**Native**  
Clump-forming species; stems have a sweet scent resembling vanilla when bruised.  
**HABITAT:** Thickets, open woods.  
**CHARACTERISTICS:**  
Height: Up to 6.1 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Pink, Purple  
Approx. Seeds Per Lb: 672,000  
Seeding Rate: Up to 1% of a mix



Herbaceous Annual/  
Biennial/Perennial

***Gaillardia pulchella***  
**ANNUAL GAILLARDIA (INDIAN BLANKET)**

**Native**  
Showy bunch-type species in annual wildflower beds.  
**HABITAT:** Dry sandy places, meadows, open areas.  
**CHARACTERISTICS:**  
Height: Up to 2 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 7.0-8.5  
Bloom Period: Spring/Fall  
Flower Color(s): Yellow  
Approx. Seeds Per Lb: 238,144  
Seeding Rate: Up to 10% of a mix



Herbaceous Biennial

**Gaura biennis**  
**BIENNIAL BEEBLOSSOM**

**Native**  
Adds diversity to meadow mixes.  
**HABITAT:** Moist meadows, stream-banks, floodplains, roadside thickets.

**CHARACTERISTICS:**  
Height: Up to 6.1 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Pink, White  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

**Helenium autumnale**  
**COMMON SNEEZEWEED**

**Native**  
Attractive late season bloomer.  
**HABITAT:** Swamps, moist river-banks, alluvial thickets, wet fields.

**CHARACTERISTICS:**  
Height: Up to 5.3 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: High  
pH: 4.0-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow  
Approx. Seeds Per Lb: 1,464,500  
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial

**Geum canadense**  
**WHITE AVENS**

**Native**  
Small seedheads add texture to landscapes after blooming.

**HABITAT:** Dry or moist woods, roadsides.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 4 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 4.5-7.5  
Bloom Period: Spring/Summer  
Flower Color(s): White  
Approx. Seeds Per Lb: 400,000  
Seeding Rate: Up to 0.5% of a mix




Herbaceous Perennial

**Helenium flexuosum**  
**PURPLEHEAD SNEEZEWEED**

**Native**  
Provides summer color to moist soils.

**HABITAT:** Wet meadows, river-banks, moist fields, roadsides.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 4.5-7.5  
Bloom Period: Spring/Fall  
Flower Color(s): Yellow  
Approx. Seeds Per Lb: 2,000,000  
Seeding Rate: Up to 1% of a mix



Herbaceous Annual

**Gypsophila elegans**  
**ANNUAL BABY'S BREATH**

**Naturalized**  
Fine-textured, fast-growing species; blooms 60 days after seeding.

**HABITAT:** Gardens, meadows.

**CHARACTERISTICS:**  
Height: Up to 2 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Bloom Period: Summer  
Flower Color(s): White  
Approx. Seeds Per Lb: 375,000  
Seeding Rate: Up to 2.5% of a mix



Herbaceous Perennial

**Helianthus angustifolius**  
**NARROWLEAF SUNFLOWER**

**Native**  
Clump-forming species; latest blooming sunflower we carry; small seeds are eaten by birds; nectar is food for migrating monarch butterflies.

**HABITAT:** Ditches, savannas, marshes, wet meadows, pine barrens.

**CHARACTERISTICS:**  
Height: Up to 6.1 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 4.0-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow  
Approx. Seeds Per Lb: 504,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Annual

***Helianthus annuus***  
**COMMON SUNFLOWER**

**Native**  
Reseeding annual; provides food for songbirds.

**HABITAT:** Meadows, roadsides.

**CHARACTERISTICS:**

Height: Up to 10.2 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.5-7.8  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 46,900  
Seeding Rate: Up to 10% of a mix



Herbaceous Perennial

***Hibiscus moscheutos***  
**CRIMSONEYED ROSEMALLOW**

**Native**  
Showy long-lived species in wet meadows and at the water's edge.

**HABITAT:** Alluvial meadows, swamp forest edges, brackish marshes.

**CHARACTERISTICS:**

Height: Up to 6.5 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
Pollinator Value: Low  
pH: 4.0-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 200,000  
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Helianthus maximiliani***  
**MAXIMILIAN'S SUNFLOWER**

**Native**  
Clump-forming species; tall shielding growth provides food and cover for birds.

**HABITAT:** Prairies, old fields, rail-road tracks, urban open ground.

**CHARACTERISTICS:**

Height: Up to 9.3 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 6.0-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 196,300  
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial

***Hypericum perforatum***  
**COMMON ST. JOHNSWORT**

**Naturalized**  
Source of herbal St. Johnswort; not for use in native mixes; seeds are persistent in soils.

**HABITAT:** Fields, roadsides, open spaces; tolerates poor soils.

**CHARACTERISTICS:**

Height: Up to 2.6 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Seeding Rate: Up to 0.1% of a mix



Herbaceous Perennial

***Heliopsis helianthoides***  
**OXEYE SUNFLOWER**

**Native**  
Vigorous clump-forming species with a long bloom period; provides food and cover for birds.

**HABITAT:** Fields, woods, floodplains, streambanks.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Spring/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 102,000  
Seeding Rate: Up to 2% of a meadow mix



Herbaceous Perennial

***Hypericum punctatum***  
**SPOTTED ST. JOHNSWORT**

**Native**  
Commonly found in riparian areas.

**HABITAT:** Moist fields, floodplains, thickets, roadsides.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: Low  
pH: 4.6-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Seeding Rate: Up to 0.2% of a mix



Herbaceous Perennial

***Hypericum pyramidatum***  
**GREAT ST. JOHNSWORT**

**Native**  
Tall showy species with large flowers.

**HABITAT:** Alluvial shores, rocky banks, swamps.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: Low  
pH: 5.7-7.1  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,800,000  
Seeding Rate: Up to 0.2% of a mix



Herbaceous Perennial  
Legume

***Lespedeza capitata***  
**ROUNDHEAD LESPEDEZA**

**Native**  
Provides food for birds and small ground animals.

**HABITAT:** Dry open woods, sand dunes, prairies.

**CHARACTERISTICS:**

Height: Up to 5.3 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
Pollinator Value: Low  
pH: 5.7-8.2  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 174,000  
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Kosteletzkya virginica***  
**VIRGINIA SALTMARSH MALLOW**

**Native**  
Attractive in coastal marshes.

**HABITAT:** Salt or brackish marshes, shores.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Pink

Approx. Seeds Per Lb: 26,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial  
Legume

***Lespedeza frutescens***  
**SHRUBBY BUSHCLOVER**


**Native**  
Clump-forming legume; one of our prettiest native Lespedezas; attractive foliage adds soft color to native meadows; seeds provide food for wildlife.

**HABITAT:** Rocky woods, thickets, dry open woods.

**CHARACTERISTICS:**

Height: Up to 3.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Low  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Seeding Rate: Up to 3% of a mix



Herbaceous Perennial  
Legume

***Lathyrus sylvestris***  
**FLAT PEA**

**Naturalized**  
Hardy rhizomatous legume; excellent for soil stabilization in infertile soils; seed must be incorporated into the soil to achieve successful establishment.

**HABITAT:** Borders of fields and thickets.

**CHARACTERISTICS:**

Height: Up to 6.5 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Shade  
Drought Tolerance: High  
pH: 5.0-7.8  
Bloom Period: Summer  
Flower Color(s): Purple, Pink, White

Approx. Seeds Per Lb: 8,000  
Seeding Rate: Up to 60% of a mix; up to 20 lb per acre alone with 15 lb per acre of tall fescue



Herbaceous Perennial  
Legume

***Lespedeza virginica***  
**SLENDER LESPEDEZA**

**Native**  
Clump-forming legume; our prettiest native Lespedeza; attractive flowers and foliage add color to native meadows; prolific producer of seeds eaten by game birds.

**HABITAT:** Open woods, roadsides, fields.

**CHARACTERISTICS:**

Height: Up to 3.8 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Low  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 175,000  
Seeding Rate: Up to 3% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

***Liatris spicata***  
**MARSH BLAZING STAR**

**Native**  
Corm-forming species; flowers have a feathery appearance.

**HABITAT:** Moist fields, fencerows, roadsides.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: High  
pH: 5.6-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 100,000  
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial  
Legume

***Lotus corniculatus***  
**BIRD'S FOOT TREFOIL**

**Naturalized**  
Excellent for controlling erosion on strip mines and landfills; provides good livestock forage.

**HABITAT:** Fields, roadsides, meadows.

**CHARACTERISTICS:**  
Height: Up to 2.3 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.0-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 369,800  
Seeding Rate: Up to 30% of a mix; up to 15 lb per acre alone



Herbaceous Perennial

***Linum perenne***  
**PERENNIAL BLUE FLAX**

**Naturalized**  
Short stature and intense blue flowers make this a great species.

**HABITAT:** Naturalized along roadsides.

**CHARACTERISTICS:**  
Height: Up to 2.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer  
Flower Color(s): Blue

Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Ludwigia alternifolia***  
**SEEDBOX**

**Native**  
Four-sided seedheads add texture to winter landscapes.

**HABITAT:** Wet woods, swampy fields.

**CHARACTERISTICS:**  
Height: Up to 3.7 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 20,800,000  
Seeding Rate: Up to 0.2% of a mix



Herbaceous Perennial

***Lobelia siphilitica***  
**GREAT BLUE LOBELIA**

**Native**  
Decorative species with indeterminate blooms.

**HABITAT:** Swamps, moist meadows, streambanks, ditches.

**CHARACTERISTICS:**  
Height: Up to 4.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Blue

Approx. Seeds Per Lb: 7,760,000  
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

***Ludwigia linearis***  
**NARROWLEAF PRIMROSE WILLOW**

**Native**  
Fast-growing species with indeterminate blooms.

**HABITAT:** Savannas, ditches, bogs.

**CHARACTERISTICS:**  
Height: Up to 2.6 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Seeding Rate: Up to 0.5% of a mix



***Ludwigia maritima***  
**SEASIDE PRIMROSE WILLOW**

**Native**  
Nice addition to a southeast coastal plain wetland mix.

**HABITAT:** Savannas, ditches, low pinelands.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Mimulus ringens***  
**SQUARE STEMMED MONKEY-FLOWER**

**Native**  
Rhizomatous species; reseeds itself in open wet areas; provides cover for wildlife; indeterminate snap-dragon-like blooms are attractive to native pollinators.

**HABITAT:** Wet open ground of swamps, meadows, shores.

**CHARACTERISTICS:**

Height: Up to 4.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 22,900,000  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Lupinus polyphyllus***  
**BIGLEAF LUPINE**

**Native**  
Very attractive in early summer; not for use in habitat restoration in our region.

**HABITAT:** Fields, roadsides; adapted to fine to coarse soils.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer  
Flower Color(s): Blue

Approx. Seeds Per Lb: 75,000  
Seeding Rate: Up to 2.5% of a mix

**Herbaceous Perennial  
Legume**



***Monarda fistulosa***  
**WILD BERGAMOT**

**Native**  
Showy rhizomatous species; often used as an ornamental.

**HABITAT:** Fields, brushy thickets, prairies, roadsides.

**CHARACTERISTICS:**

Height: Up to 4 ft.  
Minimum Root Depth: 4 in.  
Shade Tolerance: Moderate  
Drought Tolerance: No  
Pollinator Value: High  
pH: 6.0-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 1,272,500  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Lycopus americanus***  
**AMERICAN WATER HORE-  
HOUND**

**Native**  
Adds diversity to wet meadows and wetlands.

**HABITAT:** Wet ditches, swamps, moist thickets, fields, shaded hill-sides.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: High  
pH: 5.2-7.8  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 3,025,300  
Seeding Rate: Up to 0.4% of a mix

**Herbaceous Perennial**



***Monarda punctata***  
**SPOTTED BEEBALM**

**Native**  
Species with a pleasant fragrance.

**HABITAT:** Dry sandy and Coastal Plain soils, sandy upland forests, forest edges, fields, roadsides.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: High  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 1,472,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Annual/  
Biennial/Perennial**



Herbaceous Biennial

***Oenothera biennis***  
**EVENING PRIMROSE**

**Native**  
Showy the second year after seeding; new blossoms occur every evening and early morning during the season; provides good wildlife food and habitat, especially for birds.

**HABITAT:** Fields, prairies, roadsides, waste areas.

**CHARACTERISTICS:**  
Height: Up to 5.1 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 5.0-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,376,000  
Seeding Rate: Up to 0.2% of a mix



Herbaceous Annual

***Papaver rhoeas***  
**CORN POPPY**

**Naturalized**  
Creates an early field of color a year after seeding.

**HABITAT:** Roadsides, meadows, gardens.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Fall  
Flower Color(s): Red

Approx. Seeds Per Lb: 3,179,000  
Seeding Rate: Up to 0.8% of a mix



Herbaceous Perennial

***Oenothera fruticosa var. fruticosa***  
**SUNDROPS**

**Native**  
Attractive flowers.

**HABITAT:** Fields, meadows, roadsides, open woods.

**CHARACTERISTICS:**  
Height: Up to 2.9 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.5-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 3,780,000  
Seeding Rate: Up to 0.6% of a mix



Herbaceous Perennial

***Penstemon digitalis***  
**TALL WHITE BEARDTONGUE**

**Native**  
Durable, early clump-forming species; found in many of our meadow mixes.

**HABITAT:** Meadows, old fields, roadsides.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Shade  
Drought Tolerance: High  
Pollinator Value: High  
pH: 5.5-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 400,000  
Seeding Rate: Up to 4% of a mix



Herbaceous Perennial

***Oenothera speciosa***  
**SHOWY EVENING PRIMROSE**

**Native**  
Large flowers open in the evening.

**HABITAT:** Dry open places, roadsides.

**CHARACTERISTICS:**  
Height: Up to 2 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer  
Flower Color(s): White, Pink

Approx. Seeds Per Lb: 3,436,000  
Seeding Rate: Up to 0.6% of a mix



Herbaceous Perennial

***Penstemon hirsutus***  
**HAIRY BEARDTONGUE**

**Native**  
Our shortest and earliest blooming *Penstemon*; attractive addition to short grass meadows.

**HABITAT:** Dry fields, woods, roadside banks, rocky slopes.

**CHARACTERISTICS:**  
Height: Up to 2.8 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Spring/Summer  
Flower Color(s): Purple

Approx. Seeds Per Lb: 3,877,000  
Seeding Rate: Up to 0.4% of a mix



***Penstemon laevigatus***  
**APPALACHIAN BEARDTONGUE**

**Native**  
Provides early season bloom to meadows in full sun to partial shade.

**HABITAT:** Meadows, woods, roadsides.

**CHARACTERISTICS:**  
Height: Up to 3.8 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Spring/Summer  
Flower Color(s): Purple  
Seeding Rate: Up to 4.5% of a mix

**Herbaceous Perennial**



***Polygonum sagittatum***  
**ARROWLEAF TEARTHUMB**

**Native**  
Fast-growing vine; not for use in residential settings as the prickly stems can cut the skin; provides food for waterfowl.

**HABITAT:** Wet meadows, bogs, marshes.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 4.0-8.5  
Bloom Period: Spring  
Flower Color(s): Pink

Approx. Seeds Per Lb: 125,000  
Seeding Rate: Up to 2% of a mix

**Herbaceous Annual/  
Perennial Vine**



***Penthorum sedoides***  
**DITCH STONECROP**

**Native**  
Stoloniferous species; adds texture to wet landscapes; provides erosion control and habitat for wildlife.

**HABITAT:** Low wet ground, ditches.

**CHARACTERISTICS:**  
Height: Up to 2.6 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 5.0-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 45,000,000  
Seeding Rate: Up to 0.2% of a mix

**Herbaceous Perennial**



***Pycnanthemum tenuifolium***  
**NARROWLEAF MOUNTAIN-MINT**

**Native**  
Rhizomatous mint species.

**HABITAT:** Dry soils of prairies and upland woods, moist old fields, floodplains, sandy streambanks.

**CHARACTERISTICS:**  
Height: Up to 3 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 5,336,000  
Seeding Rate: Up to 0.4% of a mix

**Herbaceous Perennial**



***Polygonum pensylvanicum***  
**PENNSYLVANIA SMARTWEED**

**Native**  
Acts as a cover crop in floodplain areas and on wetland sites; provides food and cover for wildlife.

**HABITAT:** Meadows, fields, waste places, moist ditches.

**CHARACTERISTICS:**  
Height: Up to 5.7 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 4.0-8.5  
Bloom Period: Spring/Fall  
Flower Color(s): Pink

Approx. Seeds Per Lb: 126,100  
Seeding Rate: Up to 2% of a mix

**Herbaceous Annual**



***Pycnanthemum virginianum***  
**VIRGINIA MOUNTAINMINT**

**Native**  
Rhizomatous mint species; stems and leaves are fragrant when crushed; also referred to as mountain thyme.

**HABITAT:** Upland woods, moist prairies.

**CHARACTERISTICS:**  
Height: Up to 3.4 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 3,872,000  
Seeding Rate: Up to 0.4% of a mix

**Herbaceous Perennial**



2 mm Grid

Herbaceous Perennial

***Ratibida columnifera***  
**YELLOW PRAIRIE CONEFLOWER**

**Native**  
Attractive tap-rooted species in a perennial wildflower mix; source of food for birds and pollinators.

**HABITAT:** Dry open places, including prairies.

**CHARACTERISTICS:**  
Height: Up to 3.9 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.9-7.0  
Bloom Period: Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 737,100  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Rudbeckia fulgida var. fulgida***  
**ORANGE CONEFLOWER**

**Native**  
Provides decorative late summer to fall color.

**HABITAT:** Dry to moist woodlands, meadows.

**CHARACTERISTICS:**  
Height: Up to 3.4 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow, Orange

Approx. Seeds Per Lb: 500,000  
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

***Ratibida pinnata***  
**GREY HEADED CONEFLOWER**

**Native**  
Long-lasting showy flowers; provides food and cover for wildlife.

**HABITAT:** Dry prairies, dry woods, old fields.

**CHARACTERISTICS:**  
Height: Up to 4.8 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 5.6-6.8  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 427,500  
Seeding Rate: Up to 2% of a mix



Herbaceous Annual/  
Biennial/Perennial

***Rudbeckia hirta***  
**BLACKEYED SUSAN**

**Native**  
Most common native flower in our meadows; provides food and cover for birds.

**HABITAT:** Fields, meadows, roadsides.

**CHARACTERISTICS:**  
Height: Up to 3.3 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 6.0-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,575,700  
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

***Rhexia virginica***  
**VIRGINIA MEADOWBEAUTY**

**Native**  
Adds texture to fall landscapes; decorative urn-shaped seed capsules.

**HABITAT:** Bogs, ditches, low meadows.

**CHARACTERISTICS:**  
Height: Up to 3.2 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer/Fall  
Flower Color(s): Pink

Seeding Rate: Up to 0.5% of a mix



2mm Grid

Herbaceous Perennial

***Rudbeckia subtomentosa***  
**SWEET BLACKEYED SUSAN**

**Native**  
Provides food and cover for birds; nectar source for bees.

**HABITAT:** Prairies, low ground.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 712,000  
Seeding Rate: Up to 0.5% of a mix



***Rudbeckia triloba***  
**BROWNEYED SUSAN**

**Native**  
Provides late summer color and texture to landscapes; provides food for birds.

**HABITAT:** Old fields, rocky slopes, woodland edges.

**CHARACTERISTICS:**

Height: Up to 4.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow, Orange

Approx. Seeds Per Lb: 536,000  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Biennial**



***Senna marilandica***  
**MARYLAND SENNA**

**Native**  
Robust long-lived legume with sturdy growth; provides food and cover for wildlife.

**HABITAT:** Dry roadsides, rocky woodlands, streambanks.

**CHARACTERISTICS:**

Height: Up to 5.9 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 4.0-7.0  
Bloom Period: Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 20,500  
Seeding Rate: Up to 1.5% of a mix

**Herbaceous Perennial Legume**



***Saururus cernuus***  
**LIZARD'S TAIL**

**Native**  
Beautiful wetland wildflower.

**HABITAT:** Marshes, swamps.

**CHARACTERISTICS:**

Height: Up to 3.7 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer  
Flower Color(s): White

Seeding Rate: Up to 3% of a mix

**Herbaceous Perennial**



***Silphium asteriscus var. laevicaule***  
**STARRY ROSINWEED**

**Native**  
Clump-forming legume; seeds provide food for birds.

**HABITAT:** Woodlands, old fields, thickets.

**CHARACTERISTICS:**

Height: Up to 8.8 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Seeding Rate: Up to 2% of a mix

**Herbaceous Perennial**



***Senna hebecarpa***  
**WILD SENNA**

**Native**  
Robust legume with sturdy growth; provides food for birds; excellent food source for bumblebees.

**HABITAT:** Streambanks, moist old fields, moist open woods.

**CHARACTERISTICS:**

Height: Up to 6.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 20,000  
Seeding Rate: Up to 1.5% of a mix

**Herbaceous Perennial Legume**



***Silphium perfoliatum***  
**CUP PLANT**

**Native**  
Robust long-lived legume; best established by planting in the fall as a dormant seeding, with germination occurring in the spring; may be used as forage for domestic animals with multiple cuts; high potential as a bioenergy crop; provides food for birds and wildlife.

**HABITAT:** Floodplains, abandoned fields, moist meadows.

**CHARACTERISTICS:**

Height: Up to 8.5 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: High  
pH: 4.5-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 100,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



Herbaceous Perennial

***Silphium terebinthinaceum***  
**PRAIRIE DOCK**

**Native**  
Adds texture to landscapes; seeds provide food for birds.

**HABITAT:** Prairies.

**CHARACTERISTICS:**

Height: Up to 10.8 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 17,000  
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial

***Solidago canadensis***  
**CANADA GOLDENROD**

**Native**  
Aggressive rhizomatous species; does well with switchgrass and big bluestem; dominant robust vegetation adds diversity to native landscapes; provides cover for wildlife. (Note: Our harvest of this species from natural stands includes the species *Solidago gigantea* and *Solidago altissima*).

**HABITAT:** Moist or dry open places, fields, roadsides.

**CHARACTERISTICS:**

Height: Up to 6.5 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 4.8-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 4,600,000  
Seeding Rate: Up to 0.2% of a mix



Herbaceous Perennial

***Silphium trifoliatum***  
**WHORLED ROSINWEED**

**Native**  
Clump-forming species; seeds provide food for birds; source of nectar for butterflies.

**HABITAT:** Roadsides, dry fields, meadows.

**CHARACTERISTICS:**

Height: Up to 8.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 20,800  
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial

***Solidago juncea***  
**EARLY GOLDENROD**

**Native**  
First goldenrod of the season to bloom; beautiful mid-summer addition to meadows.

**HABITAT:** Open woods, fields, meadows, roadsides.

**CHARACTERISTICS:**

Height: Up to 4.2 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 2,538,000  
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

***Solidago bicolor***  
**WHITE GOLDENROD**


**Native**  
Clump-forming species; survives on low fertility sites and road cuts.

**HABITAT:** Dry woods, wooded banks, shale barrens.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 1,649,000  
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial

***Solidago nemoralis***  
**GRAY GOLDENROD**


**Native**  
Our shortest goldenrod; food source for pollinators.

**HABITAT:** Fields, open woods, roadsides in low fertility soils.

**CHARACTERISTICS:**

Height: Up to 3.7 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 6.5-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,008,000  
Seeding Rate: Up to 0.7% of a mix



***Solidago odora***  
**LICORICE SCENTED GOLDEN-ROD**


**Native**  
Showy addition to landscapes; leaves have a scent resembling black licorice when crushed.

**HABITAT:** Dry open woods, sandy soils.

**CHARACTERISTICS:**  
Height: Up to 5.4 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Seeding Rate: Up to 0.3% of a mix

**Herbaceous Perennial**



***Solidago rigida***  
**STIFF GOLDENROD**

**Native**  
Attractive rhizomatous species with robust growth; provides food and cover for wildlife.

**HABITAT:** Dry open woods, prairies.

**CHARACTERISTICS:**  
Height: Up to 5.7 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: High  
Pollinator Value: Very High  
pH: 5.0-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,009,000  
Seeding Rate: Up to 0.7% of a mix

**Herbaceous Perennial**



***Solidago patula***  
**ROUGHLEAF GOLDENROD**

**Native**  
Provides cover for wildlife.

**HABITAT:** Swamps, wet meadows, floodplains, moist woods.

**CHARACTERISTICS:**  
Height: Up to 6.5 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: Very High  
pH: 4.5-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 700,000  
Seeding Rate: Up to 1% of a mix

**Herbaceous Perennial**



***Solidago rugosa***  
**WRINKLELEAF GOLDENROD**

**Native**  
Rhizomatous species; dense stems provide cover for wildlife.

**HABITAT:** Fields, woods, floodplains, thickets, roadsides, open ground.

**CHARACTERISTICS:**  
Height: Up to 6.1 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: Very High  
pH: 5.0-7.5  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,000,000  
Seeding Rate: Up to 0.7% of a mix

**Herbaceous Perennial**



***Solidago riddellii***  
**RIDDELL'S GOLDENROD**

**Native**  
Attractive rhizomatous species; provides cover for wildlife.

**HABITAT:** Swamps, wet meadows, moist prairies.

**CHARACTERISTICS:**  
Height: Up to 4.1 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,544,000  
Seeding Rate: Up to 0.5% of a mix

**Herbaceous Perennial**



***Solidago speciosa***  
**SHOWY GOLDENROD**

**Native**  
Source of nectar for migrating monarch butterflies.

**HABITAT:** Open woods, prairies, fields, plains.

**CHARACTERISTICS:**  
Height: Up to 5.7 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,340,000  
Seeding Rate: Up to 0.6% of a mix

**Herbaceous Perennial**



**Verbena hastata**  
**BLUE VERVAIN**  
**Native**  
 Easily seeded in wetlands.  
**HABITAT:** Moist meadows, flood-plains, wet ditches, roadsides.  
**CHARACTERISTICS:**  
 Height: Up to 5.7 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: High  
 Bloom Period: Summer/Fall  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 1,488,000  
 Seeding Rate: Up to 4% of a mix

Herbaceous Biennial/  
 Perennial



**Verbesina alternifolia**  
**WINGSTEM**  
**Native**  
 Aggressive growth provides cover in wet meadows.  
**HABITAT:** Moist wooded slopes, shaded lowlands, riverbanks.  
**CHARACTERISTICS:**  
 Height: Up to 9.8 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: Very High  
 Bloom Period: Summer/Fall  
 Flower Color(s): Yellow  
 Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



**Verbena stricta**  
**HOARY VERVAIN**  
**Native**  
 Adds texture and color to open landscapes.  
**HABITAT:** Prairies, barrens, fields, roadsides; tolerates poor soils.  
**CHARACTERISTICS:**  
 Height: Up to 3.8 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: High  
 Bloom Period: Summer  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 527,000  
 Seeding Rate: Up to 2% of a mix

Herbaceous Annual/  
 Perennial



**Vernonia angustifolia**  
**TALL IRONWEED**  
**Native**  
 Showy clump-forming species.  
**HABITAT:** Savannas, pine barrens, sandy woods, old fields.  
**CHARACTERISTICS:**  
 Height: Up to 3.9 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: High  
 Bloom Period: Summer  
 Flower Color(s): Purple  
 Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



**Verbena urticifolia**  
**WHITE VERVAIN**  
**Native**  
 Adds texture to meadows.  
**HABITAT:** Thickets, moist fields, meadows, open places.  
**CHARACTERISTICS:**  
 Height: Up to 4.9 ft.  
 Pollinator Value: High  
 Bloom Period: Summer/Fall  
 Flower Color(s): White  
 Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



**Vernonia gigantea**  
**GIANT IRONWEED**  
**Native**  
 Aggressive growth provides nesting habitat for woodcock; seeds are a source of food for birds.  
**HABITAT:** Moist fields, wet woods, floodplains, meadows.  
**CHARACTERISTICS:**  
 Height: Up to 9.8 ft.  
 Shade Tolerance: Unknown  
 Drought Tolerance: Unknown  
 Pollinator Value: High  
 Bloom Period: Summer/Fall  
 Flower Color(s): Purple  
 Approx. Seeds Per Lb: 320,000  
 Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Herbaceous Perennial

***Vernonia noveboracensis***  
**NEW YORK IRONWEED**

**Native**  
Aggressive growth provides cover; seeds provide food for birds.

**HABITAT:** Streambanks, wet fields, pastures, meadows.

**CHARACTERISTICS:**

Height: Up to 6.9 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 4.5-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): Purple

Approx. Seeds Per Lb: 300,000  
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial

***Zizia aurea***  
**GOLDEN ALEXANDERS**

**Native**  
One of our earliest blooming natives; early source of food for pollinators.

**HABITAT:** Wooded bottomlands, streambanks, moist meadows, floodplains.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Shade Tolerance: Moderate  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Spring/Summer  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 172,000  
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

***Veronicastrum virginicum***  
**CULVER'S ROOT**

**Native**  
Showy rhizomatous species.

**HABITAT:** Moist meadows, thickets, swamps.

**CHARACTERISTICS:**

Height: Up to 5.9 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Summer  
Flower Color(s): Pink, White

Approx. Seeds Per Lb: 7,761,000  
Seeding Rate: Up to 0.1% of a mix



Herbaceous Annual

***Viola cornuta***  
**JOHNNY JUMPUP**

**Naturalized**  
Pansy-like miniature species; added to mixes for early spring color.

**HABITAT:** Dry meadows.

**CHARACTERISTICS:**

Height: Up to 0.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring/Summer  
Flower Color(s): White, Purple

Approx. Seeds Per Lb: 796,000  
Seeding Rate: Up to 2% of a mix



2 mm Grid

Woody Deciduous Shrub

***Amorpha canescens***  
**LEADPLANT**


**Native**  
Source of food for wildlife and pollinators.  
**HABITAT:** Sandy open woods, dry prairies.  
**CHARACTERISTICS:**  
Height: Up to 3.9 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Shade  
Drought Tolerance: High  
pH: 5.5-8.0  
Bloom Period: Summer  
Flower Color(s): Purple  
  
Approx. Seeds Per Lb: 195,000  
Seeding Rate: Up to 0.4% of a mix



Woody Deciduous Shrub

***Cephalanthus occidentalis***  
**BUTTONBUSH**

**Native**  
Produces an abundance of seed favored by ducks; good source of nectar for butterflies and bees.  
**HABITAT:** Low wet ground, swamps, bogs, streambanks, lake edges.  
**CHARACTERISTICS:**  
Height: Up to 9.8 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
pH: 4.7-8.6  
Bloom Period: Summer  
Flower Color(s): White  
  
Approx. Seeds Per Lb: 134,000  
Seeding Rate: Up to 0.6% of a wet-land mix



Woody Deciduous Shrub

***Amorpha fruticosa***  
**RIVER LOCUST**

**Native**  
Provides food and cover for wildlife; source of food for bees.  
**HABITAT:** Alluvial soils along streams, rivers, other moist areas.  
**CHARACTERISTICS:**  
Height: Up to 16.3 ft.  
Minimum Root Depth: 24 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 5.0-8.5  
Bloom Period: Spring/Summer  
Flower Color(s): Purple  
  
Approx. Seeds Per Lb: 77,000  
Seeding Rate: Up to 1% of a mix



Woody Deciduous Shrub

***Cornus amomum***  
**SILKY DOGWOOD**

**Native**  
Thick, low vegetation provides excellent habitat for wildlife; abundant fruit is eaten by birds; blue berries in the fall.  
**HABITAT:** Moist woods, fields, swamps, riparian areas.  
**CHARACTERISTICS:**  
Height: Up to 9.8 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 5.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): White  
  
Approx. Seeds Per Lb: 12,000  
Seeding Rate: Up to 6.3% of a mix



Woody Deciduous Shrub

***Amorpha herbacea***  
**CLUSTERSPIKE FALSE INDIGO**

**Native**  
Attractive plant architecture and flowers.  
**HABITAT:** Sandy fields, ridges, open woodlands; generally on the Coastal Plain.  
**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Very High  
Bloom Period: Spring/Summer  
Flower Color(s): Blue, White  
  
Seeding Rate: Up to 0.8% of a mix



Woody Deciduous Tree

***Cornus florida***  
**FLOWERING DOGWOOD**

**Native**  
Ornamental tree for woodland borders; provides abundant food for birds; red fruit.  
**HABITAT:** Woods, woodland edges.  
**CHARACTERISTICS:**  
Height: Up to 39 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 4.8-7.7  
Bloom Period: Spring/Summer  
Flower Color(s): Pink, White  
  
Approx. Seeds Per Lb: 4,500  
Seeding Rate: Up to 5.6% of a mix



***Cornus racemosa*  
GRAY DOGWOOD**

**Native**

Attractive rhizomatous shrub; thick, low vegetation provides excellent habitat for wildlife; fruit is a fall and winter food source for birds; white berries on red panicles.

**HABITAT:** Moist meadows, thickets, streambanks, roadsides.

**CHARACTERISTICS:**

Height: Up to 16.3 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Shade  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 4.8-7.4  
Bloom Period: Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 13,000  
Seeding Rate: Up to 5.8% of a mix

**Woody Deciduous Shrub**



***Ilex verticillata*  
WINTERBERRY**

**Native**

Red berries provide fall and winter food for wildlife.

**HABITAT:** Swamps, bogs, moist woods, wet shores.

**CHARACTERISTICS:**

Height: Up to 16.3 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Pollinator Value: Low  
pH: 4.5-7.5  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 92,000  
Seeding Rate: Up to 0.8% of a mix

**Woody Deciduous Shrub**



***Cornus sericea*  
RED OSIER DOGWOOD**

**Native**

Attractive stoloniferous shrub; thick, low vegetation provides excellent habitat for wildlife; bright red stems add winter color; abundant white fruit is eaten by birds.

**HABITAT:** Moist woods, moist meadows, thickets, riparian areas, swamps.

**CHARACTERISTICS:**

Height: Up to 9.8 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 4.8-7.5  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 18,000  
Seeding Rate: Up to 4.2% of a mix

**Woody Deciduous Shrub**



***Parthenocissus quinquefolia*  
VIRGINIA CREEPER**

**Native**

High-growing vine; produces black fruit in the fall.

**HABITAT:** Woods, fields, woodland edges.

**CHARACTERISTICS:**

Height: Up to 49 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Moderate  
Drought Tolerance: High  
Pollinator Value: Low  
pH: 5.0-7.5  
Bloom Period: Spring/Summer  
Flower Color(s): Green

Approx. Seeds Per Lb: 18,000  
Seeding Rate: Up to 1% of a mix

**Woody Deciduous Vine**



***Hamamelis virginiana*  
WITCHHAZEL**

**Native**

Provides mid-story habitat; nuts provide fall food for wildlife.

**HABITAT:** Moist woods, brushy fields.

**CHARACTERISTICS:**

Height: Up to 16.3 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.5-6.2  
Bloom Period: Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 11,000  
Seeding Rate: Up to 6.8% of a mix

**Woody Deciduous Shrub**



***Platanus occidentalis*  
AMERICAN SYCAMORE**

**Native**

Large riverbank tree; provides shade and cover for riparian shrubs and forbs.

**HABITAT:** Streambanks, low woods, floodplains.

**CHARACTERISTICS:**

Height: Up to 162.5 ft.  
Minimum Root Depth: 30 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 4.9-6.5  
Bloom Period: Spring/Summer

Approx. Seeds Per Lb: 192,000  
Seeding Rate: Up to 0.1% of a mix

**Woody Deciduous Tree**



Woody Deciduous Tree

**Prunus serotina**  
**WILD BLACK CHERRY**

**Native**  
One of North America's most valuable hardwoods; provides food (fruit) and nesting sites for wildlife; source of food for caterpillars of 300 species of moths and butterflies eaten by birds; purplish-black berries in the fall.

**HABITAT:** Woods, fencerows.

**CHARACTERISTICS:**

Height: Up to 97.5 ft.  
Minimum Root Depth: 36 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 4.0-7.5  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 4,800  
Seeding Rate: Up to 5.2% of a mix



Woody Deciduous Tree

**Robinia pseudoacacia**  
**BLACK LOCUST**

**Native**  
Provides nitrogen and cover for successive vegetation; good honey producer.

**HABITAT:** Reclamation sites, floodplains, thickets, fencerows; develops quickly in poor soils.

**CHARACTERISTICS:**

Height: Up to 97.5 ft.  
Minimum Root Depth: 36 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 4.6-8.2  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 24,000  
Seeding Rate: Up to 1% of a mix



Woody Deciduous Shrub

**Prunus virginiana**  
**CHOCHECHERRY**

**Native**  
Creates intermediate to mid-story habitat for nesting; abundant fruit provides summer and fall food for birds; host to many native caterpillars that are also food for birds; dark red to purple berries in the summer and fall.

**HABITAT:** Rocky upland woods, roadsides.

**CHARACTERISTICS:**

Height: Up to 32.5 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: High  
pH: 5.2-8.4  
Bloom Period: Spring/Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 5,000  
Seeding Rate: Up to 5% of a mix



Woody Deciduous Shrub

**Rosa carolina**  
**PASTURE ROSE**

**Native**  
Provides cover for wildlife.

**HABITAT:** Dunes, prairies, fields, upland woods.

**CHARACTERISTICS:**

Height: Up to 6.5 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: High  
Pollinator Value: Medium  
pH: 4.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Pink

Approx. Seeds Per Lb: 50,000  
Seeding Rate: Up to 1.5% of a mix



Woody Deciduous Shrub

**Rhus typhina**  
**STAGHORN SUMAC**

**Native**  
Bark provides winter food for cottontail rabbits; fruit is an emergency winter food source for birds, including wild turkey.

**HABITAT:** Dry open fields, roadsides, woodland edges.

**CHARACTERISTICS:**

Height: Up to 32.5 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Summer  
Flower Color(s): Green

Approx. Seeds Per Lb: 60,000  
Seeding Rate: Up to 1.3% of a mix



Woody Deciduous Shrub

**Rosa palustris**  
**SWAMP ROSE**

**Native**  
Attractive rhizomatous shrub; provides food and cover for songbirds, deer, ruffed grouse, pheasants, and wild turkey; develops rose hips that last into winter.

**HABITAT:** Streambanks, wetlands, marshes.

**CHARACTERISTICS:**

Height: Up to 8.1 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 4.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Pink

Seeding Rate: Up to 3% of a mix



Woody Deciduous Tree

***Salix amygdaloides*  
PEACHLEAF WILLOW**

**Native**  
Good bioengineering material; fibrous roots grow into the water table to stabilize the streambank; leaf drops replenish organic life in streams; provides shade and habitat for wildlife.

**HABITAT:** Riparian areas, wetlands.

**CHARACTERISTICS:**

Height: Up to 65 ft.  
Minimum Root Depth: 30 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 6.0-8.0  
Bloom Period: Spring

Seeding Rate: Space live stakes on 3'-6' centers; may be used as larger live post plantings



Woody Deciduous Shrub

***Salix exigua ssp. interior*  
SANDBAR WILLOW**

**Native**  
Excellent rhizomatous bioengineering material; tolerant of ice and debris loading from streamflow; provides habitat for wildlife.

**HABITAT:** Sandbars, sandy or gravel streambanks, waterways.

**CHARACTERISTICS:**

Height: Up to 19.5 ft.  
Minimum Root Depth: 36 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.0-7.8  
Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Tree

***Salix discolor*  
PUSSY WILLOW**

**Native**  
Good bioengineering material; early show of catkins is the first sign of spring; provides habitat for wildlife.

**HABITAT:** Wetlands, moist or wet woods.

**CHARACTERISTICS:**

Height: Up to 22.8 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.0-7.0  
Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Shrub

***Salix lucida*  
SHINING WILLOW**

**Native**  
Good bioengineering material; well-suited for wetland restoration; provides streambank erosion protection and habitat for wildlife.

**HABITAT:** Wetlands, streambanks.

**CHARACTERISTICS:**

Height: Up to 19.5 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.8-7.2  
Bloom Period: Spring  
Flower Color(s): Yellow

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Shrub

***Salix eriocephala*  
HEARTLEAF WILLOW**

**Native**  
Good rhizomatous bioengineering material; provides habitat for wildlife.

**HABITAT:** Riparian areas, streambanks.

**CHARACTERISTICS:**

Height: Up to 19.5 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
pH: 4.0-7.0  
Bloom Period: Spring/Summer

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Tree

***Salix nigra*  
BLACK WILLOW**

**Native**  
Stems can be brittle, making this only a fair bioengineering material; leaf drops replenish organic life in streams; provides shade and habitat for wildlife.

**HABITAT:** Wet meadows, riparian areas.

**CHARACTERISTICS:**

Height: Up to 65 ft.  
Minimum Root Depth: 32 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.8-8.0  
Bloom Period: Spring/Summer

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Shrub

***Salix purpurea***  
**STREAMCO WILLOW**

**Naturalized**  
Excellent early rooting bioengineering material; leaf drops replenish organic life in streams; provides shade to small streams and habitat for wildlife.

**HABITAT:** Streambanks, riparian areas.

**CHARACTERISTICS:**  
Height: Up to 16.3 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: No  
pH: 5.5-7.5  
Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Shrub

***Sambucus canadensis***  
**ELDERBERRY**

**Native**  
Purplish-black berries provide an excellent source of summer food for wildlife.

**HABITAT:** Woods, moist fields, streambanks, moist roadsides.

**CHARACTERISTICS:**  
Height: Up to 13 ft.  
Minimum Root Depth: 16 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
Pollinator Value: Medium  
pH: 5.0-8.9  
Bloom Period: Summer  
Flower Color(s): White, Yellow

Approx. Seeds Per Lb: 292,000  
Seeding Rate: Up to 0.3% of a mix



Woody Deciduous Shrub

***Salix sericea***  
**SILKY WILLOW**

**Native**  
Excellent bioengineering material; provides dense habitat for wildlife.

**HABITAT:** Wetlands, streambanks, riparian areas.

**CHARACTERISTICS:**  
Height: Up to 13 ft.  
Minimum Root Depth: 18 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.2-7.0  
Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers



Woody Deciduous Shrub

***Viburnum dentatum***  
**ARROWWOOD**

**Native**  
Provides late summer and fall food, cover, browse, and nesting sites for birds; bluish-black fruit.

**HABITAT:** Wet woods, swamps.

**CHARACTERISTICS:**  
Height: Up to 16.3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Pollinator Value: Medium  
Bloom Period: Spring/Summer  
Flower Color(s): Pink, White

Approx. Seeds Per Lb: 20,000  
Seeding Rate: Up to 3.8% of a mix; space rooted seedlings on 6' centers



Woody Deciduous Shrub


***Salix x cottetii***  
**DWARF WILLOW**

**Naturalized**  
Excellent bioengineering material; ideal for utility stream crossings; tolerant of ice and debris loading from streamflow; resilient species that will recover from vehicle disturbance; provides good habitat for wildlife.

**HABITAT:** Streambanks.

**CHARACTERISTICS:**  
Height: Up to 8 ft.  
Minimum Root Depth: 20 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.5-7.5  
Bloom Period: Spring

Seeding Rate: Space live stakes on 2' centers



Woody Deciduous Shrub

***Viburnum lentago***  
**NANNYBERRY**

**Native**  
Dense foliage provides cover; bluish-black fruit is a food source for wildlife.

**HABITAT:** Woods, wetlands, roadsides.

**CHARACTERISTICS:**  
Height: Up to 32.5 ft.  
Minimum Root Depth: 14 in.  
Shade Tolerance: Shade  
Drought Tolerance: Low  
Pollinator Value: Medium  
pH: 5.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Pink, White

Approx. Seeds Per Lb: 7,843  
Seeding Rate: Up to 9.6% of a mix



**Woody Deciduous Shrub**

***Viburnum trilobum***  
**AMERICAN CRANBERRY**

**Native**

Red berries provide early winter food for wildlife.

**HABITAT:** Wetlands, wet woods.

**CHARACTERISTICS:**

Height: Up to 16.3 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Full Sun

Drought Tolerance: No

Pollinator Value: Medium

pH: 5.5-7.5

Bloom Period: Spring/Summer

Flower Color(s): White

Approx. Seeds Per Lb: 13,600

Seeding Rate: Up to 5.5% of a mix



**2 mm Grid**

**Herbaceous Perennial**

***Arrhenatherum elatius ssp. elatius***  
**TALL OATGRASS, 'RUFFNER'**

**Naturalized**  
 Bunchgrass with a pronounced cool season growth habit; used for forage and as a cover crop; good into Canada; highly palatable for whitetail deer throughout late fall and early winter.

**HABITAT:** Roadsides, fields, waste ground; persists in shallow, moderately infertile soils.

**CHARACTERISTICS:**  
 Height: Up to 6.5 ft.  
 Minimum Root Depth: 14 in.  
 Shade Tolerance: Moderate  
 Drought Tolerance: Moderate  
 pH: 5.0-7.0  
 Bloom Period: Summer

Approx. Seeds Per Lb: 189,000  
 Seeding Rate: Up to 10 lb per acre when direct drilled; up to 20 lb per acre when broadcast seeded



**Herbaceous Annual**

***Brassica napus***  
**FORAGE RAPE, 'BONAR'**

**Naturalized**  
 Late maturing, high yielding forage rape; blooms in the spring if planted in the fall; blooms 70 days post-planting if planted in the spring; quality winter feed for cattle and sheep; palatable for all livestock and grazing wildlife.

**HABITAT:** Pastures; grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**  
 Height: Up to 2.3 ft.  
 Minimum Root Depth: 6 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Low  
 pH: 6.0-7.2  
 Bloom Period: Spring/Summer

Seeding Rate: Up to 5 lb per acre when direct drilled



**Herbaceous Annual**


***Avena sativa***  
**OATS**

**Naturalized**  
 Small, bunch-type cereal grain for human and animal consumption; companion cover crop with a spring or fall seeding; matures quickly in hot weather and killed by freezing winter weather.

**HABITAT:** Tolerates a wide range of soil types, but prefers fertile open areas.

**CHARACTERISTICS:**  
 Height: Up to 4 ft.  
 Minimum Root Depth: 8 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Low  
 pH: 5.3-8.5  
 Bloom Period: Summer

Approx. Seeds Per Lb: 19,000  
 Seeding Rate: Up to 90 lb per acre as a grain crop; up to 30 lb per acre as a cover crop with a perennial planting



**2 mm Grid**

**Herbaceous Annual**


***Brassica napus***  
**FORAGE RAPE, 'RANGIORA'**

**Naturalized**  
 Late maturing, high yielding forage rape; blooms in the spring if planted in the fall; blooms 70 days post-planting if planted in the spring; quality winter feed for cattle and sheep; palatable for all livestock and grazing wildlife.

**HABITAT:** Pastures; grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**  
 Height: Up to 2.3 ft.  
 Minimum Root Depth: 6 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Low  
 pH: 6.0-7.2  
 Bloom Period: Spring/Summer

Seeding Rate: Up to 5 lb per acre when direct drilled



**2 mm Grid**

**Herbaceous Annual/Biennial**

***Brassica napus***  
**FORAGE BRASSICA, 'WINFRED'**

**Naturalized**  
 Blooms in the spring if planted in the fall; blooms 70 days post-planting if planted in the spring; leafy growth is palatable for deer, cattle, and sheep.

**HABITAT:** Food plots, forage (pastures)

**CHARACTERISTICS:**  
 Height: Up to 1.8 ft.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Low  
 pH: 5.5-7.5  
 Bloom Period: Spring  
 Flower Color(s): Yellow

Approx. Seeds Per Lb: 157,000  
 Seeding Rate: Up to 8 lb per acre when direct drilled



**Herbaceous Annual/Biennial**


***Brassica napus***  
**WINTER CANOLA**

**Naturalized**  
 Can be very productive if planted in the fall; seeds are approximately 37% oil; also used in the production of healthy cooking oils; high quality livestock supplement; preferred by deer in the winter; makes good biodiesel.

**HABITAT:** Agricultural soils that support winter wheat production; grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**  
 Height: Up to 6 ft.  
 Minimum Root Depth: 6 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Low  
 pH: 6.0-7.2  
 Bloom Period: Spring  
 Flower Color(s): Yellow

Approx. Seeds Per Lb: 160,000  
 Seeding Rate: Up to 10 lb per acre when direct drilled



2 mm Grid

Herbaceous Annual/Biennial

**Brassica napus**  
**DWARF ESSEX RAPE**

**Naturalized**


Member of the mustard family; seeds are approximately 37% oil; used in the production of healthy cooking oils; high quality livestock supplement; preferred by deer in the winter; makes good biodiesel.

**HABITAT:** Agricultural soils that support winter wheat production; grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**

Height: Up to 6 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
pH: 6.0-7.2  
Bloom Period: Spring  
Flower Color(s): Yellow

Seeding Rate: Up to 10 lb per acre when direct drilled



2 mm Grid

Herbaceous Annual/Biennial

**Brassica rapa**  
**HYBRID FORAGE TURNIP,**  
**'PACER'**

**Naturalized**

Tap-rooted cross between forage rape and forage turnip; high energy, early maturing hybrid forage brassica; excellent regrowth (50-70 days); excellent for grazing.

**HABITAT:** Grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**

Height: Up to 2 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Bloom Period: Spring

Seeding Rate: Up to 5 lb per acre when direct drilled



2 mm Grid

Herbaceous Annual/Biennial

**Brassica rapa**  
**HYBRID FORAGE BRASSICA,**  
**'VIVANT'**

**Naturalized**


Tap-rooted brassica; used by dairy cattle, beef cattle, and sheep; graze within 35-50 days of planting and at 25-30-day intervals thereafter.

**HABITAT:** Grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**

Height: Up to 2 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Bloom Period: Spring

Seeding Rate: Up to 6 lb per acre when direct drilled



2 mm Grid

Herbaceous Annual/Biennial

**Brassica rapa**  
**PURPLE TOP TURNIP**

**Naturalized**

Tap-rooted species; leaves are high in protein; highly palatable and digestible for deer.

**HABITAT:** Grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Bloom Period: Spring

Seeding Rate: Up to 4 lb per acre when direct drilled



Herbaceous Annual

**Brassica rapa**  
**FORAGE TURNIP, 'APPIN'**

**Naturalized**


Also referred to as field mustard; establishes in 50-80 days; 80% digestible and high in protein and carbohydrates; tolerates multiple grazings; multi-crowned for improved regrowth.

**HABITAT:** Pastures, fields, fertile open areas; grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
Bloom Period: Spring

Approx. Seeds Per Lb: 192,800  
Seeding Rate: Up to 5 lb per acre when direct drilled



2 mm Grid

Herbaceous Annual/Biennial

**Brassica spp.**  
**BRASSICA, FORAGE, 'RANGI'**

**Naturalized**

Provides food for wildlife.

**HABITAT:** Fields, waste ground.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.0-8.0  
Bloom Period: Spring

Seeding Rate: Up to 5 lb per acre when direct drilled



Herbaceous Annual/Biennial

**Camelina sativa**  
**CAMELINA**

**Naturalized**

Member of the mustard family; used in the production of culinary oils and biodiesel; very low requirements for fertility, tillage, and weed control; oil and meal are high in Omega 3 fatty acids; approved by the USDA for human, poultry, and livestock consumption.

**HABITAT:** Prospers in various climates and soils; well-suited for marginal soils.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Summer  
Flower Color(s): White, Yellow

Approx. Seeds Per Lb: 400,000  
Seeding Rate: Up to 6 lb per acre when direct drilled



Herbaceous Annual

**Fagopyrum esculentum**  
**BUCKWHEAT**

**Naturalized**

Tap-rooted, renovation grain cover crop for low productivity land; builds organic matter that decays rapidly when plowed under; planting time varies with application; killed by frost; provides food for human consumption, wildlife, and honey production.

**HABITAT:** Adapts to a wide range of soil types and conditions.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.0-8.5  
Bloom Period: Summer

Approx. Seeds Per Lb: 20,000  
Seeding Rate: Up to 75 lb per acre alone (not recommended as a companion crop)



Herbaceous Perennial

**Cichorium spp.**  
**CHICORY**

**Naturalized**

High-yielding tap-rooted species; not for ornamental purposes; selected for intense grazing, less bolting, and better disease resistance; 20% protein; suitable for all wildlife and production livestock; excellent for-age quality.

**HABITAT:** Grows best in moderately drained, medium to high fertility soils.

**CHARACTERISTICS:**

Height: Up to 2 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: High  
pH: 5.6-6.5  
Bloom Period: Summer

Seeding Rate: Up to 2 lb per acre in a mix; up to 5 lb per acre alone



Herbaceous Perennial

**Festuca elatior x Lolium perenne**  
**FESTULOLIUM, 'DUO'**

**Naturalized**

Endophyte-free, high sugar cross between meadow fescue and a tetraploid perennial ryegrass; excellent palatability.

**HABITAT:** Fertile soils.

**CHARACTERISTICS:**

Height: Up to 2.7 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Moderate  
Bloom Period: Summer

Seeding Rate: Up to 25 lb per acre in a mix; up to 30 lb per acre alone



Herbaceous Annual

**Echinochloa crusgalli var. frumentacea**  
**JAPANESE MILLET**

**Naturalized**

Warm season bunchgrass; used for erosion control and as a fast-growing summer companion crop; seed in the spring or summer; after growth, may be flooded to a depth of 18" during waterfowl migration season; provides food for wildlife and a favorite of waterfowl.

**HABITAT:** Well-drained soils, but thrives in wetlands.

**CHARACTERISTICS:**

Height: Up to 4.9 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 4.7-7.4  
Bloom Period: Summer

Approx. Seeds Per Lb: 143,000  
Seeding Rate: Up to 30 lb per acre as a grain crop; up to 10 lb per acre as a companion crop



Herbaceous Annual Legume

**Glycine max**  
**SOYBEAN, 'BOBWHITE'**

**Naturalized**

Trailing legume for use in wildlife food plots; matures 120 days after planting.

**HABITAT:** Well to moderately drained soils and full sun.

**CHARACTERISTICS:**

Height: Up to 4 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 6.5-7.5  
Bloom Period: Summer  
Flower Color(s): Purple

Seeding Rate: Up to 60 lb per acre alone



***Helianthus spp.***  
**BLACK OIL SUNFLOWER**

**Naturalized**  
 Grain is used for oil and meal; meal provides quality livestock feed; extremely popular food source for birds; may be left in the field for winter bird food.

**HABITAT:** Food plots.


**CHARACTERISTICS:**

Height: Up to 9.8 ft.  
 Minimum Root Depth: 8 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Moderate  
 pH: 6.0  
 Bloom Period: Summer  
 Flower Color(s): Yellow

Approx. Seeds Per Lb: 7,000  
 Seeding Rate: Up to 10% of a food plot mix; up to 20 lb per acre alone or 0.5 lb per 1,000 sq ft

2 mm Grid

Herbaceous Annual



***Melilotus officinalis***  
**YELLOW BLOSSOM SWEET-CLOVER**

**Naturalized**  
 Tap-rooted legume; builds soil organic matter on low fertility sites; source of nectar and pollen for honeybees.

**HABITAT:** Mine spoil, roadsides, waste places.

**CHARACTERISTICS:**

Height: Up to 6.5 ft.  
 Minimum Root Depth: 32 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: High  
 pH: 5.0-8.0  
 Bloom Period: Summer  
 Flower Color(s): Yellow

Approx. Seeds Per Lb: 259,000  
 Seeding Rate: Up to 5 lb per acre in a mix; up to 20 lb per acre alone

Herbaceous Biennial Legume



***Medicago sativa***  
**ALFALFA**

**Naturalized**  
 Hardy component of right-of-way mixes; grows quickly after mowing; good livestock and wildlife forage.

**HABITAT:** Well-drained high fertility soils.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
 Minimum Root Depth: 24 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: High  
 pH: 6.0-8.5  
 Bloom Period: Spring  
 Flower Color(s): Purple

Approx. Seeds Per Lb: 227,000  
 Seeding Rate: Up to 10 lb per acre in a mix; up to 20 lb per acre alone

Herbaceous Perennial Legume



***Onobrychis viciifolia***  
**SAINFOIN**

**Naturalized**  
 Provides forage for deer, elk, cattle, and sheep; do not graze for two seasons following planting.

**HABITAT:** Hay fields, wildlife food plots; less persistent in moist soils.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
 Minimum Root Depth: 14 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: High  
 pH: 6.0-8.5  
 Bloom Period: Spring/Summer  
 Flower Color(s): Red

Approx. Seeds Per Lb: 30,000  
 Seeding Rate: Up to 25% of a mix; up to 34 lb per acre alone

2 mm Grid

Herbaceous Perennial Legume



***Melilotus alba***  
**WHITE BLOSSOM SWEETCLOVER**

**Naturalized**  
 Tap-rooted legume; source of nectar and pollen for bees.

**HABITAT:** Roadsides, waste places.

**CHARACTERISTICS:**

Height: Up to 9.8 ft.  
 Minimum Root Depth: 32 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: High  
 Bloom Period: Summer/Fall  
 Flower Color(s): White

Approx. Seeds Per Lb: 259,000  
 Seeding Rate: Up to 5 lb per acre in a mix; up to 20 lb per acre alone

Herbaceous Biennial Legume



***Panicum miliaceum***  
**WHITE PROSO MILLET**

**Naturalized**  
 Grain for human and livestock consumption; seeds are one of the most attractive foods for birds.

**HABITAT:** Well-drained loamy soils; tolerates various soil conditions.

**CHARACTERISTICS:**

Height: Up to 3.3 ft.  
 Minimum Root Depth: 10 in.  
 Shade Tolerance: Full Sun  
 Drought Tolerance: Low  
 pH: 5.8-6.8  
 Bloom Period: Summer

Seeding Rate: Up to 35% of a mix; up to 20 lb per acre alone

2 mm Grid

Herbaceous Annual



Herbaceous Perennial

**Phleum pratense**  
**TIMOTHY**

**Naturalized**  
Bunch-type grass; excellent forage for horses and other livestock.

**HABITAT:** Fields, open areas; moderate fertility requirements.

**CHARACTERISTICS:**  
Height: Up to 4.9 ft.  
Minimum Root Depth: 10 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 5.5-7.0  
Bloom Period: Summer/Fall  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 1,230,000  
Seeding Rate: Up to 90% of a mix; up to 10 lb per acre alone



Herbaceous Perennial

**Sanguisorba minor**  
**SMALL BURNET**

**Naturalized**  
Very attractive to deer for its herbal taste.

**HABITAT:** Sunny flatlands to open slopes in well-drained soils.

**CHARACTERISTICS:**  
Height: Up to 2.3 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Moderate  
Drought Tolerance: Low  
pH: 6.0-8.0  
Bloom Period: Spring/Summer  
Flower Color(s): Red

Approx. Seeds Per Lb: 48,700  
Seeding Rate: Up to 10% of a food plot mix; up to 2 lb per acre with clover



Herbaceous Annual Legume

**Pisum arvense**  
**AUSTRIAN WINTER PEA**

**Naturalized**  
Cold-tolerant cool season legume; used for erosion control as a cover crop or temporary hay crop; builds nitrogen and organic matter in fields and gardens; not winter hardy north of the Mason-Dixon Line; seed in early spring or fall; excellent for wild game food plots.

**HABITAT:** Prefers dry soils.

**CHARACTERISTICS:**  
Height: Up to 4 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring

Seeding Rate: Up to 50% of a mix; up to 60 lb per acre alone



Herbaceous Annual

**Secale cereale**  
**RYE**

**Naturalized**  
Bunch-type winter companion or cover crop; used for erosion control; may be planted anytime of year, but preferably in the fall or winter as rye has a strong ability to grow in cold weather.

**HABITAT:** More productive than other cereals in infertile sandy or acidic soils.

**CHARACTERISTICS:**  
Height: Up to 5 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 4.5-8.2  
Bloom Period: Spring  
Flower Color(s): Yellow

Approx. Seeds Per Lb: 18,000  
Seeding Rate: Up to 90 lb per acre as a grain crop; up to 30 lb per acre as a cover crop with a perennial planting



Herbaceous Annual


**Raphanus sativus**  
**RADISH, 'GROUNDHOG'**

**Naturalized**  
Species with a taproot that can reach a depth of 6"; root exudates help to suppress nematodes; plant at least 60 days prior to frost.

**HABITAT:** Cultivated; occasionally escapes to roadsides or old fields.

**CHARACTERISTICS:**  
Height: Up to 3 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Unknown  
Bloom Period: Spring  
Flower Color(s): Pink, Purple, White

Seeding Rate: Up to 10 lb per acre drilled with up to 30 lb per acre of grain rye



Herbaceous Annual

**Setaria italica**  
**GERMAN MILLET**

**Naturalized**  
Warm season bunchgrass; used as a cover crop and for pasture and haylage; good supplemental hay for cattle and sheep; seed in the spring or summer; provides food for wildlife.

**HABITAT:** Grows best in well-drained loamy soils with low moisture.

**CHARACTERISTICS:**  
Height: Up to 5 ft.  
Minimum Root Depth: 8 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.3-6.9  
Bloom Period: Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 217,000  
Seeding Rate: Up to 30 lb per acre as a grain crop; up to 10 lb per acre as a cover crop



2 mm Grid

Herbaceous Annual

**Sorghum spp.**  
**SORGHUM/RED MILO**

**Naturalized**

Early maturing species; most attractive to mourning doves in the fall; makes emergency food for turkey, pheasants, and deer during the winter.

**HABITAT:** Cultivated fields.

**CHARACTERISTICS:**

Height: Up to 5 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: High  
pH: 5.5-7.5  
Bloom Period: Summer

Approx. Seeds Per Lb: 27,000  
Seeding Rate: Up to 50% of a mix; up to 30 lb per acre with a grain drill or when broadcast seeded



Herbaceous Annual Legume

**Trifolium incarnatum**  
**CRIMSON CLOVER**

**Naturalized**

Used as a winter companion or cover crop in pasture, hay, and silage mixes and for erosion control; seed in the fall or early spring south of I-64; plow under before the next crop is planted; provides food for honeybees.

**HABITAT:** Sandy and clay-like soils.

**CHARACTERISTICS:**

Height: Up to 2.6 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 6.0-7.0  
Bloom Period: Spring/Summer  
Flower Color(s): Red

Approx. Seeds Per Lb: 150,000  
Seeding Rate: Up to 10 lb per acre as a companion crop with a wildflower mix; up to 30 lb per acre alone



2 mm Grid

Herbaceous Annual Legume

**Trifolium alexandrinum**  
**BERSEEM CLOVER**

**Naturalized**


Forage legume for deer and livestock.

**HABITAT:** Hay fields, food plots.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 6.5-8.0  
Bloom Period: Summer  
Flower Color(s): White

Approx. Seeds Per Lb: 207,000  
Seeding Rate: Up to 20% in a mix with alfalfa; up to 14 lb per acre drill seeded



2 mm Grid

Herbaceous Annual Legume

**Trifolium michelianum**  
**BALANSA CLOVER, 'FIXATION'**

**Naturalized**

Legume with a deep taproot that helps to break up soil compaction; provides food for livestock and deer.

**HABITAT:** Hay fields, food plots.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Shade Tolerance: Unknown  
Drought Tolerance: Unknown  
Bloom Period: Spring  
Flower Color(s): White

Seeding Rate: Up to 20% of a mix; up to 8 lb per acre as a monoculture



Herbaceous Perennial Legume

**Trifolium hybridum**  
**ALSIKE CLOVER**

**Naturalized**

Winter-hardy legume; provides forage for domestic animals and food and cover for wildlife.

**HABITAT:** Meadows, disturbed areas; tolerates low fertility.

**CHARACTERISTICS:**

Height: Up to 4 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.6-7.5  
Bloom Period: Spring/Summer  
Flower Color(s): Pink, White

Approx. Seeds Per Lb: 680,000  
Seeding Rate: Up to 25% of a mix; up to 10 lb per acre alone



Herbaceous Perennial Legume

**Trifolium pratense**  
**RED CLOVER**

**Naturalized**

Short-lived legume; used as hay, pasture, and silage for domestic animals, or as a cover crop that builds nitrogen and organic matter; flowers develop again after early summer cutting; may be frost seeded during early spring; provides food for bumblebees.

**HABITAT:** Medium fertility soils.

**CHARACTERISTICS:**

Height: Up to 2.6 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 6.0-7.6  
Bloom Period: Spring/Summer  
Flower Color(s): Pink

Approx. Seeds Per Lb: 272,000  
Seeding Rate: Up to 25% of a mix; up to 10 lb per acre alone



**Herbaceous Perennial  
Legume**

*Trifolium repens*  
**WHITE CLOVER**

**Naturalized**

Good stoloniferous erosion control cover crop; grows again after mowing or grazing; highly palatable as food for wildlife and domestic animals; source of food for honeybees.

**HABITAT:** Moist soils, lawns, field borders.

**CHARACTERISTICS:**

Height: Up to 1 ft.  
Minimum Root Depth: 12 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.2-8.0  
Bloom Period: Summer/Fall  
Flower Color(s): White

Approx. Seeds Per Lb: 712,000  
Seeding Rate: Up to 10% of a mix; up to 10 lb per acre alone



**Herbaceous Annual**

*Urochloa ramosa*  
**BROWN TOP MILLET**

**Naturalized**

Short-lived species for pasture, hay, or wildlife forage; establishes quickly for erosion control; plant from May to August; matures in 60 days.

**HABITAT:** Tolerates low fertility acidic soils.

**CHARACTERISTICS:**

Height: Up to 3 ft.  
Minimum Root Depth: 24 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Low  
pH: 5.5-6.9  
Bloom Period: Summer

Approx. Seeds Per Lb: 75,000  
Seeding Rate: Up to 20 lb per acre when direct drilled; up to 30 lb per acre when broadcast seeded; up to 10 lb per acre as a cover crop



**Herbaceous Biennial Legume**

*Vicia villosa*  
**HAIRY VETCH**

**Naturalized**

Builds nitrogen and organic matter in fields and gardens; seed in early fall after row crop is harvested; likely to freeze at -30 degrees F.

**HABITAT:** Well-drained soils.

**CHARACTERISTICS:**

Height: Up to 7 ft.  
Minimum Root Depth: 6 in.  
Shade Tolerance: Full Sun  
Drought Tolerance: Moderate  
pH: 6.0-7.5  
Bloom Period: Spring  
Flower Color(s): Purple

Approx. Seeds Per Lb: 16,000  
Seeding Rate: Up to 10 lb per acre in a mix with up to 50 lb per acre of grain rye

INDEX

INDEX TO SCIENTIFIC/COMMON NAMES

NAME	PAGE	NAME	PAGE	NAME	PAGE
<i>Achillea millefolium</i>	85	Hairy	101	<i>lupulina</i>	73
<i>Acorus americanus</i>	69	Tall White	101	<i>lurida</i>	73
<i>Agastache foeniculum</i>	85	Beebalm, Spotted	100	<i>pensylvanica</i>	73
<i>Agrimonia parviflora</i>	85	Beeblossom, Biennial	96	<i>scoparia</i>	73
Agrimony, Small Flowered	85	Beggartick	89	<i>shortiana</i>	73
<i>Agrostis</i>		Bentgrass		<i>stipata</i>	73
<i>alba</i>	69	Autumn	69	<i>stricta</i>	74
<i>hyemalis</i>	69	Creeping	69	<i>vulpinoidea</i>	74
<i>perennans</i>	69	Colonial	69	<i>Centaurea cyanus</i>	90
<i>stolonifera</i>	69	Winter	69	<i>Cephalanthus occidentalis</i>	109
<i>tenuis</i>	69	Bergamot, Wild	100	<i>Chamaecrista</i>	
Alfalfa	118	<i>Bidens</i>		<i>fasciculata</i>	90
<i>Alisma subcordatum</i>	70	<i>aristosa</i>	89	<i>nictitans</i>	90
Alkaligrass	81	<i>cernua</i>	89	<i>Chasmanthium</i>	
<i>Allium cernuum</i>	70	<i>frondosa</i>	89	<i>latifolium</i>	74
<i>Alopecurus arundinaceus</i>	70	Biennial Beeblossom	96	<i>laxum</i>	74
<i>Amorpha</i>		Black Cherry, Wild	111	<i>Cheiranthus allionii</i>	90
<i>canescens</i>	109	Blackeyed Susan	103	<i>Chelone glabra</i>	90
<i>fruticosa</i>	109	Sweet	103	Chicory	117
<i>herbacea</i>	109	Blanketflower	95	Blue	91
<i>Amsonia ciliata</i>	85	Blazing Star, Marsh	99	Chokecherry	111
<i>Andropogon</i>		Blue Eyed Grass, Narrowleaf	82	<i>Chrysanthemum</i>	
<i>gerardii</i>	70	Blueflag	78	<i>leucanthemum</i>	90
<i>ternarius</i>	70	Bluegrass		<i>maximum</i>	91
<i>virginicus</i>	70	Fowl	80	Chufa	74
<i>Anemone</i>		Kentucky	80	<i>Cichorium</i>	
<i>canadensis</i>	85	Rough	81	<i>intybus</i>	91
Canadian	85	Bluejoint, Canada	71	<i>spp.</i>	117
<i>virginiana</i>	85	Bluestar, Fringed	85	<i>Cinna arundinacea</i>	74
<i>Anthoxanthum odoratum</i>	71	Bluestem		<i>Clematis virginiana</i>	91
<i>Apocynum cannabinum</i>	86	Big	70	Clover	
<i>Aquilegia canadensis</i>	86	Little	81	Alsike	120
<i>Arrhenatherum elatius ssp. elatius</i>	115	Splitbeard	70	Balansa, 'Fixation'	120
Arrow Arum	80	Boneset	95	Berseem	120
Arrowwood	113	Bottlebrush Grass	75	Crimson	120
<i>Asclepias</i>		<i>Bouteloua</i>		Red	120
<i>incarnata</i>	86	<i>curtipendula</i>	71	White	121
<i>syriaca</i>	86	<i>gracilis</i>	71	Columbine, Eastern	86
<i>tuberosa</i>	86	<i>Brassica</i>		Coneflower	
<i>Aster</i>		Forage, 'Rangi'	116	Grey Headed	103
Aromatic	87	Forage, 'Winfred'	115	Orange	103
Arrowleaf	88	Hybrid Forage, 'Vivant'	116	Pale Purple	94
Bigleaf	87	<i>napus</i>	115, 116	Purple	94
Calico	87	<i>rapa</i>	116	Yellow Prairie	103
<i>divaricatus</i>	86	<i>spp.</i>	116	Cordgrass, Prairie	83
Flat Topped White	88	Brome, Smooth	71	<i>Coreopsis</i>	
Heath	87	<i>Bromus inermis</i>	71	<i>basalis</i>	91
<i>laevis</i>	87	Broomsedge	70	<i>grandiflora</i>	91
<i>lateriflorus</i>	87	Browneyed Susan	104	Lanceleaf	91
<i>macrophyllus</i>	87	Buckwheat	117	<i>lanceolata</i>	91
New England	87	Bulrush		Plains	92
<i>novae-angliae</i>	87	Green	81	Tall	92
<i>oblongifolius</i>	87	Northern	82	<i>tinctoria</i>	92
<i>pilosus</i>	87	Softstem	82	<i>tripteris</i>	92
<i>preanthoides</i>	88	Wood	82	Cornflower	90
<i>puniceus</i>	88	Bundleflower, Illinois	93	<i>Cornus</i>	
Purplestem	88	Bur Marigold	89	<i>amomum</i>	109
<i>sagittifolius</i>	88	Nodding	89	<i>florida</i>	109
Showy	88	Bur Reed		<i>racemosa</i>	110
Smooth Blue	87	Eastern	83	<i>sericea</i>	110
<i>spectabilis</i>	88	Giant	83	<i>Coronilla varia</i>	92
<i>umbellatus</i>	88	Burnet, Small	119	Cosmos	92
White Wood	86	Bushclover, Shrubby	98	<i>bipinnatus</i>	92
Zigzag	88	Buttonbush	109	Sulphur	92
<i>Avena sativa</i>	115	<i>Calamagrostis canadensis</i>	71	<i>sulphureus</i>	92
Avens, White	96	Camelina	117	Cranberry, American	114
Baby's Breath, Annual	96	<i>Camelina sativa</i>	117	Crownvetch	92
Bachelor's Button	90	Canola, Winter	115	Cup Plant	104
<i>Baptisia</i>		<i>Carex</i>		Cutgrass, Rice	78
<i>alba</i>	88	<i>albolutescens</i>	71	Culver's Root	108
<i>australis</i>	89	<i>comosa</i>	72	<i>Cyperus esculentus</i>	74
<i>pendula</i>	89	<i>crinita</i>	72	<i>Dactylis glomerata</i>	75
<i>tinctoria</i>	89	<i>frankii</i>	72	Daisy	
Barnyardgrass	75	<i>granularis var. haleana</i>	72	Oxeye	90
Beardtongue		<i>grayi</i>	72	Shasta	91
Appalachian	102	<i>intumescens</i>	72	<i>Dalea</i>	

INDEX TO SCIENTIFIC/COMMON NAMES

NAME	PAGE	NAME	PAGE	NAME	PAGE
<i>candida</i>	92	Annual	95	<i>virginica</i>	98
<i>purpurea</i>	93	<i>aristata</i>	95	<i>Liatris spicata</i>	99
<i>Daucus carota</i>	93	Perennial	95	<i>Linum perenne</i>	99
Deertongue	79	<i>pulchella</i>	95	Lizard's Tail	104
<i>Delphinium ajacis</i>	93	Gamagrass, Eastern	84	<i>Lobelia</i>	
<i>Deschampsia cespitosa</i>	75	<i>Gaura biennis</i>	96	Great Blue	99
<i>Desmanthus illinoensis</i>	93	<i>Geum canadense</i>	96	<i>siphilitica</i>	99
<i>Desmodium</i>		<i>Glyceria</i>		Locust	
<i>canadense</i>	93	<i>grandis</i>	77	Black	111
<i>paniculatum</i>	93	<i>striata</i>	78	River	109
<i>Dianthus barbatus</i>	94	<i>Glycine max</i>	117	<i>Lolium</i>	
Dock, Prairie	105	Golden Alexanders	108	<i>multiflorum</i>	79
Dogwood		Goldenrod		<i>perenne</i>	79
Flowering	109	Canada	105	<i>Lotus corniculatus</i>	99
Gray	110	Early	105	Lovegrass	
Red Osier	110	Grassleaf	95	Bigtop	76
Silky	109	Gray	105	Purple	76
Dropseed		Licorice Scented	106	Weeping	76
Prairie	83	Riddell's	106	<i>Ludwigia</i>	
Rough	83	Roughleaf	106	<i>alternifolia</i>	99
Sand	83	Showy	106	<i>linearis</i>	99
Duck Potato	81	Stiff	106	<i>maritima</i>	100
<i>Echinacea</i>		White	105	Lupine, Bigleaf	100
<i>pallida</i>	94	Wrinkleleaf	106	<i>Lupinus polyphyllus</i>	100
<i>purpurea</i>	94	Gramma		<i>Lycopus americanus</i>	100
<i>Echinochloa</i>		Blue	71	Mannagrass	
<i>crusgalli</i> var. <i>frumentacea</i>	117	Sideoats	71	American	77
<i>muricata</i>	75	<i>Gypsophila elegans</i>	96	Fowl	78
Elderberry	113	Hairgrass, Tufted	75	Meadowbeauty, Virginia	103
<i>Elymus</i>		<i>Hamamelis virginiana</i>	110	<i>Medicago sativa</i>	118
<i>canadensis</i>	75	<i>Helenium</i>		<i>Melilotus</i>	
<i>glabriflorus</i>	75	<i>autumnale</i>	96	<i>alba</i>	118
<i>hystrix</i>	75	<i>flexuosum</i>	96	<i>officinalis</i>	118
<i>riparius</i>	76	<i>Helianthus</i>		Milkweed	
<i>virginicus</i>	76	<i>angustifolius</i>	96	Butterfly	86
<i>Eragrostis</i>		<i>annuus</i>	97	Common	86
<i>curvula</i>	76	<i>maximiliani</i>	97	Swamp	86
<i>hirsuta</i>	76	<i>spp.</i>	118	Millet	
<i>spectabilis</i>	76	<i>Heliopsis helianthoides</i>	97	Brown Top	121
<i>Eryngium yuccifolium</i>	94	<i>Hibiscus moscheutos</i>	97	German	119
<i>Eupatorium</i>		Horehound, American Water	100	Japanese	117
<i>coelestinum</i>	94	<i>Hypericum</i>		White Proso	118
<i>fistulosum</i>	94	<i>perforatum</i>	97	Milo, Red	120
<i>maculatum</i>	95	<i>punctatum</i>	97	<i>Mimulus ringens</i>	100
<i>perfoliatum</i>	95	<i>pyramidatum</i>	98	Mistflower	94
<i>purpureum</i>	95	Hyssop		<i>Monarda</i>	
<i>Euthamia graminifolia</i>	95	Anise	85	<i>fistulosa</i>	100
Evening Primrose	101	Lavender	85	<i>punctata</i>	100
Showy	101	<i>Ilex verticillata</i>	110	Monkeyflower, Square Stemmed	100
<i>Fagopyrum esculentum</i>	117	Indian Blanket	95	Mountainmint	
False Indigo		Indiangrass	82	Narrowleaf	102
Blue	89	Indianhemp	86	Virginia	102
Clusterspike	109	<i>Iris versicolor</i>	78	Nannyberry	113
Yellow	89	Ironweed		Oatgrass, Tall, 'Ruffner'	115
Fern, Sensitive	79	Giant	107	Oats	115
Fescue		New York	108	River	74
Blue	77	Tall	107	<i>Oenothera</i>	
Chewings	77	Joe Pye Weed	94	<i>biennis</i>	101
Creeping Red	77	Purple Node	95	<i>fruticosa</i> var. <i>fruticosa</i>	101
Hard	77	Spotted	95	<i>speciosa</i>	101
Sheep	77	Johnny Jumpup	108	Onion, Nodding	70
Tall	76	<i>Juncus</i>		<i>Onobrychis viciifolia</i>	118
<i>Festuca</i>		<i>effusus</i>	78	<i>Onoclea sensibilis</i>	79
<i>arundinacea</i>	76	<i>tenuis</i>	78	Orchardgrass	75
<i>elatiior x Lolium perenne</i>	117	Junegrass	78	Panicgrass	
<i>ovina</i>	77	<i>Koeleria macrantha</i>	78	Beaked	79
<i>ovina</i> var. <i>duriuscula</i>	77	<i>Kosteletzkya virginica</i>	98	Coastal	79
<i>ovina</i> var. <i>glauca</i>	77	Larkspur, Rocket	93	Redtop	80
<i>rubra</i>	77	<i>Lathyrus sylvestris</i>	98	Smooth	80
<i>rubra</i> ssp. <i>commutata</i>	77	Leadplant	109	<i>Panicum</i>	
Festulolium, 'Duo'	117	<i>Leersia oryzoides</i>	78	<i>amarum</i>	79
Flat Pea	98	<i>Lespedeza</i>		<i>anceps</i>	79
Flax, Perennial Blue	99	<i>capitata</i>	98	<i>clandestinum</i>	79
Foxtail, Creeping	70	<i>frutescens</i>	98	<i>dichotomiflorum</i>	80
<i>Gaillardia</i>		Roundhead	98	<i>miliaceum</i>	118
		Slender	98	<i>rigidulum</i>	80

INDEX TO SCIENTIFIC/COMMON NAMES

NAME	PAGE	NAME	PAGE	NAME	PAGE
<i>virgatum</i>	80	Ryegrass		<i>americanum</i>	83
<i>Papaver rhoeas</i>	101	Annual	79	<i>eurycarpum</i>	83
<i>Parthenocissus quinquefolia</i>	110	Perennial	79	<i>Spartina pectinata</i>	83
Pea		<i>Sagittaria latifolia</i>	81	Spiderwort	
Austrian Winter	119	Sainfoin	118	Ohio	84
Partridge	90	<i>Salix</i>		Virginia	84
Sensitive	90	<i>amygdaloides</i>	112	Zigzag	84
<i>Peltandra virginica</i>	80	<i>discolor</i>	112	<i>Sporobolus</i>	
<i>Penstemon</i>		<i>eriocephala</i>	112	<i>asper</i>	83
<i>digitalis</i>	101	<i>exigua ssp. interior</i>	112	<i>cryptandrus</i>	83
<i>hirsutus</i>	101	<i>lucida</i>	112	<i>heterolepis</i>	83
<i>laevigatus</i>	102	<i>nigra</i>	112	St. Johnswort	
<i>Penthorum sedoides</i>	102	<i>purpurea</i>	113	Common	97
<i>Phleum pratense</i>	119	<i>sericea</i>	113	Great	98
Pickerelweed	81	<i>x cottetii</i>	113	Spotted	97
<i>Pisum arvense</i>	119	Saltmarsh Mallow, Virginia	98	Stonecrop, Ditch	102
Plantain, Mud	70	<i>Sambucus canadensis</i>	113	Sumac, Staghorn	111
<i>Platanus occidentalis</i>	110	<i>Sanguisorba minor</i>	119	Sundrops	101
<i>Poa</i>		<i>Saururus cernuus</i>	104	Sunflower	
<i>palustris</i>	80	<i>Schizachyrium scoparium</i>	81	Black Oil	118
<i>pratensis</i>	80	<i>Scirpus</i>		Common	97
<i>trivialis</i>	81	<i>atrovirens</i>	81	Maximilian's	97
<i>Polygonum</i>		<i>cyperinus</i>	82	Narrowleaf	96
<i>pensylvanicum</i>	102	<i>expansus</i>	82	Oxeye	97
<i>sagittatum</i>	102	<i>hattorianus</i>	82	Sweetclover	
<i>Pontederia cordata</i>	81	<i>validus</i>	82	White Blossom	118
Poppy, Corn	101	<i>Secale cereale</i>	119	Yellow Blossom	118
Prairie Clover		Sedge		Sweetflag	69
Purple	93	Awl	73	Sweetwilliam	94
White	92	Blunt Broom	73	Switchgrass	80
Primrose Willow		Cosmos	72	Sycamore, American	110
Narrowleaf	99	Fox	74	Tearthumb, Arrowleaf	102
Seaside	100	Frank's	72	Thimbleweed	85
<i>Prunus</i>		Fringed	72	Tickseed	
<i>serotina</i>	111	Gray's	72	Goldenmane	91
<i>virginiana</i>	111	Greenwhite	71	Largeflower	91
<i>Puccinellia distans</i>	81	Hop	73	Tickseed Sunflower, Showy	89
Purpletop	84	Limestone Meadow	72	Ticktrefoil	
<i>Pycnanthemum</i>		Lurid	73	Panicleleaf	93
<i>tenuifolium</i>	102	Pennsylvania	73	Showy	93
<i>virginianum</i>	102	Short's	73	Timothy	119
Queen Anne's Lace	93	Star	72	<i>Tradescantia</i>	
Radish, 'GroundHog'	119	Tussock	74	<i>ohiensis</i>	84
Rape		Seedbox	99	<i>subaspera</i>	84
Dwarf Essex	116	<i>Senna</i>		<i>virginiana</i>	84
Forage, 'Bonar'	115	<i>hebecarpa</i>	104	Trefoil, Bird's Foot	99
Forage, 'Rangiora'	115	<i>marilandica</i>	104	<i>Tridens flavus</i>	84
<i>Raphanus sativus</i>	119	Maryland	104	<i>Trifolium</i>	
<i>Ratibida</i>		Wild	104	<i>alexandrinum</i>	120
<i>columnifera</i>	103	<i>Setaria italica</i>	119	<i>hybridum</i>	120
<i>pinnata</i>	103	<i>Silphium</i>		<i>incarnatum</i>	120
Rattlesnake Master	94	<i>asteriscus var. laevicaule</i>	104	<i>micelianum</i>	120
Redtop	69	<i>perfoliatum</i>	104	<i>pratense</i>	120
Reedgrass, Wood	74	<i>terebinthaceum</i>	105	<i>repens</i>	121
<i>Rhexia virginica</i>	103	<i>trifoliatum</i>	105	<i>Tripsacum dactyloides</i>	84
<i>Rhus typhina</i>	111	<i>Sisyrinchium angustifolium</i>	82	Turnip	
<i>Robinia pseudoacacia</i>	111	Smartweed, Pennsylvania	102	Forage, 'Appin'	116
<i>Rosa</i>		Sneezeweed		Hybrid Forage, 'Pacer'	116
<i>carolina</i>	111	Common	96	Purple Top	116
<i>palustris</i>	111	Purplehead	96	Turtlehead	90
Rose		<i>Solidago</i>		<i>Urochloa ramosa</i>	121
Pasture	111	<i>bicolor</i>	105	<i>Verbena</i>	
Swamp	111	<i>canadensis</i>	105	<i>hastata</i>	107
Rosemallow, Crimsoyed	97	<i>juncea</i>	105	<i>stricta</i>	107
Rosinweed		<i>nemoralis</i>	105	<i>urticifolia</i>	107
Starry	104	<i>odora</i>	106	<i>Verbesina alternifolia</i>	107
Whorled	105	<i>patula</i>	106	Vernalgrass, Sweet	71
<i>Rudbeckia</i>		<i>riddellii</i>	106	<i>Vernonia</i>	
<i>fulgida var. fulgida</i>	103	<i>rigida</i>	106	<i>angustifolia</i>	107
<i>hirta</i>	103	<i>rugosa</i>	106	<i>gigantea</i>	107
<i>subtomentosa</i>	103	<i>speciosa</i>	106	<i>noveboracensis</i>	108
<i>triloba</i>	104	<i>Sorghastrum nutans</i>	82	<i>Veronicastrum virginicum</i>	108
Rush		Sorghum	120	Vervain	
Path	78	<i>spp.</i>	120	Blue	107
Soft	78	Soybean, 'Bobwhite'	117	Hoary	107
Rye	119	<i>Sparganium</i>		White	107

INDEX

INDEX TO SCIENTIFIC/COMMON NAMES

NAME	PAGE
Vetch, Hairy	121
<i>Viburnum</i>	
<i>dentatum</i>	113
<i>lentago</i>	113
<i>trilobum</i>	114
<i>Vicia villosa</i>	121
<i>Viola cornuta</i>	108
Virgin's Bower	91
Virginia Creeper	110
Wallflower	90
Wild Indigo	
Largeleaf	89
White	88
Wildrice	84
Wildrye	
Canada	75
Riverbank	76
Southeastern	75
Virginia	76
Willow	
Black	112
Dwarf	113
Heartleaf	112
Peachleaf	112
Pussy	112
Sandbar	112
Shining	112
Silky	113
Streamco	113
Wingstem	107
Winterberry	110
Witchhazel	110
Woodoats, Slender	74
Woolgrass	82
Yarrow, Common	85
<i>Zizania aquatica</i>	84
<i>Zizia aurea</i>	108

## ACKNOWLEDGMENTS

### Some information in this catalog courtesy of:

- Bell, C. Ritchie and Taylor, Bryan J. *Florida Wild Flowers and Roadside Plants*. Laurel Hill Press, 1982.
- Brown, Melvin L. and Brown, Russell G. *Herbaceous Plants of Maryland*. Port City Press, Inc., 1984.
- Brown, Russell G. and Brown, Melvin L. *Woody Plants of Maryland*. Port City Press, 1972.
- Clewell, Andre F. *Guide to the Vascular Plants of the Florida Panhandle*. Florida State University Press, 1985.
- Duncan, Wilbur H. and Kartesz, John T. *Vascular Flora of Georgia - An Annotated Checklist*. University of Georgia Press, 1981.
- Flora of North America Editorial Committee. *Flora of North America Association*. Oxford University Press, 2006.
- Gleason Ph.D., Henry A. and Cronquist, Ph.D., *Arthur. Manual of Vascular Plants of Northeastern United States and Adjacent Canada*. 2nd ed. New York Botanical Garden, 1991.
- Magee, Dennis W. and Ahles, Harry E. *Flora of the Northeast: A Manual of the Vascular Flora of New England and Adjacent New York*. University of Massachusetts Press, 1999.
- Mohlenbrock, R.H. *Guide to the Vascular Flora of Illinois*. Southern Illinois University Press, 1986.
- Radford, Albert E., Ahles, Harry E. and Bell, C. Ritchie. *Manual of the Vascular Flora of the Carolinas*. University of North Carolina Press, 1964.
- Rhoads, Ann. Fowler & Block, Timothy A. *The Plants of Pennsylvania*. University of Pennsylvania Press, 2000.
- Rhoads, Ann Fowler and Klein, Jr., William McKinley. *The Vascular Flora of Pennsylvania: Annotated Checklist and Atlas*. American Philosophical Society, 1993.
- Richardson, L.L.; Adler, L.S.; Andiococha, J.; Regan, K.H.; Anthony, W.; Manson, J.S.; and Irwin, R.E. *Secondary metabolites in floral nectar reduce parasite infections in bumblebees*. *Proceedings of the Royal Society B*. 282:20142471,2015.
- Taylor, Walter Kingsley. *Florida Wildflowers in Their Natural Communities*. University Press of Florida, 1998.
- Taylor, Walter Kingsley. *The Guide to Florida Wildflowers*. Taylor Trade Publishing, 1992.
- The Great Plains Flora Association. *Flora of the Great Plains*. University Press of Kansas, 1986.
- Tobe, Dr. John D.; Burks, Kathy Craddock; Cantrell, Richard W.; Garland, Mark A.; Sweeley, Maynard E.; Hall, Dr. David W.; Wallace, Pete; Anglin, Guy; Nelson, Gil; Cooper, Dr. James R.; Bickner, David; Gilbert, Katherine, Aymond, Neil; Greenwood, Ken; and Raymond, Nina. *Florida Wetland Plants - An Identification Manual*. University of Florida, 1998.
- USDA, NRCS (2015). *The PLANTS Database* (<http://plants.usda.gov>, 23 October 2015). National Plant Data Team, Greensboro, NC 27401-4901 USA.
- Weakley, Alan S. *Flora of the Southern and Mid-Atlantic States*. University of North Carolina Herbarium, Working draft of 30 November 2012.
- Weakley, Alan S., Ludwig, J. Christopher and Townsend, John F. 2012. *Flora of Virginia*. Bland Crowder ed. Foundation of the Flora of Virginia Project, Inc., Richmond. Fort Worth: Botanical Research Institute of Texas Press.
- Wunderlin, Richard P. *Guide to the Vascular Plants of Central Florida*. University Press of Florida, 1982.
- Yatskievych, K. *Field Guide to Indiana Wildflowers*. Indiana University Press, 2000

### Ernst Conservation Seeds would like to thank:

- The Xerces Society for Invertebrate Research and Anne Stine, Ph.D. for providing the pollinator ratings used in this catalog.
- Harland Patch, Ph.D., Assistant Research Professor, Pennsylvania State University, Dept. of Agricultural Sciences; Research Associate, Entomology at Penn State's Center for Pollinator Research.
- Dave Biddinger, Ph.D., Tree Fruit Research Entomologist and Research Professor, Pennsylvania State University, Fruit Research & Extension Center.
- Missouri Botanical Garden Plant Finder Tool.

### Photographs in this catalog contributed by the following Ernst Seeds' employees:

Andy Ernst  
 Dan Arnett  
 Marcia Ernst  
 Mark Fiely  
 Ayshea Heckman  
 Greg Kedzierski  
 Nikki Hindle

Will Adams, Allison Walters Daisley, Sammie Boyer, Ben Crawl, Luke Fiely, Shayna Lantz, Sydney Winter, and Aarick Zook for assistance with data collection.

Photos contributed by outside sources are gratefully acknowledged on images or in captions throughout the catalog.

We're proud of the reputation we've built and value the relationships we've cultivated with our customers for more than half a century.

We look forward to serving your ecological seed needs for many years to come.

ERNSTSEED.COM  
800-873-3321  
MEADVILLE, PENNSYLVANIA

